

Climate Change and Fisheries Research in the Chesapeake Bay

A compilation of fisheries-based climate
research and tools compiled by the
NOAA Chesapeake Bay Office

November 2016



This compilation of fisheries-based climate research and tools was compiled by the NOAA Chesapeake Bay Office. The contents of this document were compiled based on a review of both peer-reviewed and grey literature as related to the 2014 Chesapeake Bay Agreement Goal to “increase the resiliency of the Chesapeake Bay Watershed, including its living resources, habitats, public infrastructure and communities, to withstand adverse impacts from changing environmental and climate conditions.” This document is not intended to be an exhaustive list; recognizing that there are additional research and tool development under way within the region. This document will be updated on an annual basis.

Please contact Zoë Johnson (zoe.johnson@noaa.gov) to suggest content for the next edition of this publication.

Table of Contents

	Page Number
<u>Section I: Oysters</u>	4
<u>Section II: Blue Crab</u>	4
<u>Section III: Striped Bass</u>	5
<u>Section IV: Forage</u>	5
<u>Section V: Submerged Aquatic Vegetation</u>	6
<u>Section VI: “Finfish” Non-specific</u>	6
<u>Section VII: General Biota</u>	7
<u>Section VIII: Habitat</u>	8
<u>Section IX: Tools</u>	10

Section I: Oysters

Kimmel, D.G., Tarnowski, M., Newell, R.I.E. 2012. Long-term (1939 to 2008) spatial patterns in juvenile eastern oyster (*Crassostrea virginica*, Gmelin 1791) abundance in the Maryland portion of Chesapeake Bay. *Journal of Shellfish Research* 31, 1023-1031.

Levinton, J., Doall, M., Ralston, D., Starke, A., Allam, B. 2011. Climate change, precipitation and impacts on an estuarine refuge from disease. *PLoS ONE* 6(4): e18849.
doi:10.1371/journal.pone.0018849.

Waldbusser, G.G., Steenson, R.A., Green, M.A. 2011. Oyster shell dissolution rates in estuarine waters: effects of pH and shell legacy. *Journal of Shellfish Research* 30, 659-669.

Waldbusser, G., Voigt, E., Bergschneider, H., Green, M., Newell, R. 2011. Biocalcification in the Eastern Oyster (*Crassostrea virginica*) in relation to long-term trends in Chesapeake Bay pH. *Estuaries and Coasts* 34(2), 221-231.

Kimmel, D.G. and Newell, R.I. 2007. The influence of climate variation on eastern oyster (*Crassostrea virginica*) juvenile abundance in Chesapeake Bay. *Limnology and Oceanography*, 52(3), pp.959-965.

Cook, T., Folli, M., Klinck, J., Ford, S. and Miller, J. 1998. The relationship between increasing sea-surface temperature and the northward spread of *Perkinsus marinus* (Dermo) disease epizootics in oysters. *Estuarine, Coastal and Shelf Science*, 46(4), pp.587-597.

Ford, S.E. 1996. Range extension by the oyster parasite *Perkinsus marinus* into the northeastern United States: response to climate change? *Oceanographic Literature Review*, 12(43), p.1265.

Section II: Blue Crab

Eggleston, D.B., N.B. Reynolds, L.L. Etherington, G.R. Plaia, and L. Xie. 2010. Tropical storm and environmental forcing on regional blue crab (*Callinectes sapidus*) settlement. *Fisheries Oceanography* 19:89-106.

Hines, A.H., Johnson, E.G., Darnell, M.Z., Rittschof, D., Miller, T.J., Bauer, L.J., Rodgers, P., Aguilar, R. 2010. Predicting effects of climate change on blue crabs in Chesapeake Bay. In: G.H. Kruse, G.L. Eckert, R.J. Foy, R.N. Lipcius, B. Sainte-Marie, D.L. Stram, D. Woodby (Editors), *Biology and Management of Exploited Crab Populations under Climate Change*. Alaska Sea Grant, University of Alaska Fairbanks, pp. 109-127.

Bauer, L., and T. Miller. 2006. Winter mortality of juvenile blue crabs in Chesapeake Bay. Masters Thesis, University of Maryland.

Aguilar, R., A.H. Hines, T.G. Wolcott, D.L. Wolcott, M.A. Kramer, and R.N. Lipcius. 2005. The timing and route of movement and migration of post-copulatory female blue crabs, *Callinectes sapidus* Rathbun, from the upper Chesapeake Bay. *Journal of Experimental Marine Biology and Ecology* 319:117-128.

Section III: Striped Bass

Martino, E.J. and Houde, E.D. 2010. Recruitment of striped bass in Chesapeake Bay: spatial and temporal environmental variability and availability of zooplankton prey. *Marine Ecology Progress Series*, 409, pp.213-228.

Coutant, C.C. 1990. Temperature-oxygen habitat for freshwater and coastal striped bass in a changing climate. *Transactions of the American Fisheries Society*, 119(2), pp.240-253.

Coutant, C.C. 1985. Striped bass, temperature, and dissolved oxygen: a speculative hypothesis for environmental risk. *Transactions of the American Fisheries Society*, 114(1), pp.31-61.

Price, K. S., D. A. Flemer, J. Taft, G. B. Mackiernan, W. Nehlsen, R. B. Biggs, N. Burger, and D. A. Blaylock. 1985. Nutrient enrichment of Chesapeake Bay and its impact on the habitat of Striped Bass: A speculative hypothesis, *Trans. Amer. Fish. Soc.*, 114(1): 97–106.

Section IV: Forage

Buchheister, A., Houde, E.D. 2016. Forage indicators and consumption profiles for Chesapeake Bay fishes, Final Report. University of Maryland Center for Environmental Science, Chesapeake Biological Laboratory, Ref. No. [UMCES] CBL 2016-001.

Miller SH, Breitburg DL, Burrell RB, Keppel AG. 2016. Acidification increases sensitivity to hypoxia in important forage fishes. *Mar Ecol Prog Ser* 549:1-8.

Prosser, D. 2014. Chesapeake Bay Forage Base: Waterbird Predation and Population Trends. United States Geological Survey Patuxent Wildlife Research Center.

Section V: Submerged Aquatic Vegetation (SAV)

Zimmerman, R.C., Hill, V.J. and Gallegos, C.L. 2015. Predicting effects of ocean warming, acidification, and water quality on Chesapeake region eelgrass. *Limnology and Oceanography*, 60(5), pp.1781-1804.

Jarvis, J.C., Brush, M.J., Moore, K.A. 2014. Modeling loss and recovery of *Zostera marina* beds in the Chesapeake Bay: The role of seedlings and seed-bank viability. *Aquatic Botany* 113, 32-45.

Orth, R.J., Williams, M.R., Marion, S., Wilcox, D.J., Carruthers, T.J.B., Moore, K.A., Kemp, W.M., Dennison, W.C., Rybicki, N., Bergstrom, P., Batiuk, R. 2010. Long-term trends in submersed aquatic vegetation (SAV) in Chesapeake Bay, USA, related to water quality. *Estuaries and Coasts* 33, 1144–1163.

Moore, K.A. and Jarvis, J.C. 2008. Environmental factors affecting recent summertime eelgrass diebacks in the lower Chesapeake Bay: implications for long-term persistence. *Journal of Coastal Research*, pp.135-147.

Palacios, S.L. and Zimmerman, R.C. 2007. Response of eelgrass *Zostera marina* to CO₂ enrichment: possible impacts of climate change and potential for remediation of coastal habitats. *Marine Ecology-Progress Series*, 344, p.1.

Section VI: “Finfish” Non-specific

Jones, C.M. 2013. Can we predict the future: juvenile finfish and their seagrass nurseries in the Chesapeake Bay. *ICES Journal of Marine Science* doi:10.1093/icesjms/fst142.

Nye, J.A., Link, J.S., Hare, J.A, and W.J. Overholtz. 2009. Changing spatial distribution of fish stocks in relation to climate and population size on the Northeast United States continental shelf, *Marine Ecology Progress*, 393:111-129.

Wood, R. J., D. F. Boesch, and V. S. Kennedy. 2002. Future consequences of climate change for the Chesapeake Bay ecosystem and its fisheries, *Am. Fish. Soc. Symp.*, 32: 171–184.

Wood, R. J. 2000. Synoptic scale climatic forcing of multispecies fish recruitment patterns in Chesapeake Bay, Ph.D. thesis, University of Virginia, School of Marine Science, The College of William and Mary, Gloucester Point, VA.

Rombough, P.J. 1997, January. The effects of temperature on embryonic and larval development. In *Seminar Series-Society For Experimental Biology* (Vol. 61, pp. 177-224). Cambridge University Press.

Frank, K.T., Perry, R.I. and Drinkwater, K.F. 1990. Predicted response of Northwest Atlantic invertebrate and fish stocks to CO₂-induced climate change. *Transactions of the American Fisheries Society*, 119(2), pp.353-365.

Section VII: General Biota

Hare JA, Morrison WE, Nelson MW, Stachura MM, Teeters EJ, et al. 2016. A Vulnerability Assessment of Fish and Invertebrates to Climate Change on the Northeast U.S. Continental Shelf. *PLoS ONE* 11(2): e0146756. doi: 10.1371/journal.pone.0146756.

Vaquier-Sunyer, R., Duarte, C.M. 2011. Temperature effects on oxygen thresholds for hypoxia in marine benthic organisms. *Global Change Biology* 17, 1788-1797.

Johnson Z.P. et. al., "Bay and Aquatic Ecosystems." Comprehensive Strategy for Reducing Maryland's Vulnerability to Climate Change, Phase II: Building Societal, Economic, and Ecological Resilience. A report to the Maryland Commission on Climate Change from the Adaptation and Response Working Group. 2010. Boicourt KE and Johnson ZP (eds). University of Maryland Center for Environmental Science, Cambridge, Maryland and Maryland Department of Natural Resources, Annapolis, Maryland.

Najjar, R.G., Pyke, C.R., Adams, M.B., Breitburg, D., Hershner, C., Kemp, M., Howarth, R., Mulholland, M.R., Paolisso, M., Secor, D. and Sellner, K. 2010. Potential climate-change impacts on the Chesapeake Bay. *Estuarine, Coastal and Shelf Science*, 86(1), pp.1-20.

Codispoti, L.A. 2010. Interesting times for marine N₂O. *Science*, 327(5971), pp.1339-1340.

Stranko S.A., R.H. Hilderbrand, R.P. Morgan III, M.W. Staley, A.J. Becker, A. Roseberry-Lincoln, E.S. Perry, and P.T. Jacobson. 2008. Brook trout declines with changes to land cover and temperature in Maryland. *North American Journal of Fisheries Management* 28: 1223-1232.

U. S. Environmental Protection Agency. 2008. Climate Change and Aquatic Invasive Species, Washington, D.C., EPA/600/R-08/014.

Pyke, C.R., Najjar, R.G., Adams, M.B., Breitburg, D., Kemp, M., Hershner, C., Howarth, R., Mulholland, M., Paolisso, M., Secor, D. and Sellner, K. 2008. Climate Change and the Chesapeake Bay: State-of-the-Science Review and Recommendations. A Report from the Chesapeake Bay Program Science and Technical Advisory Committee (STAC). Annapolis, MD, 59.

Boesch, D.F., Coles, V.J., Kimmel, D.G. and Miller, W.D. 2007. Coastal dead zones and global climate change: ramifications of climate change for Chesapeake Bay hypoxia. *Regional impacts of climate change: four case studies in the United States*, pp.57-70.

Twilley, R.R., Kleypas, J.A., Cowan Jr, J.H. and Hare, S.R. 2002. Coastal and marine ecosystems & Global climate change: Potential Effects on US Resources (pp. 1-64). Arlington, VA: Pew Center on Global Climate Change.

Wood, R.J., Boesch, D.F. and Kennedy, V.S. 2002. Future consequences of climate change for the Chesapeake Bay ecosystem and its fisheries. In *American Fisheries Society Symposium* (Vol. 32, pp. 171-184).

Najjar, R.G., Walker, H.A., Anderson, P.J., Barron, E.J., Bord, R.J., Gibson, J.R., Kennedy, V.S., Knight, C.G., Megonigal, J.P., O'Connor, R.E. and Polsky, C.D. 2000. The potential impacts of climate change on the mid-Atlantic coastal region. *Climate Research*, 14(3), pp.219-233.

Breitburg, D.L., Loher, T., Pacey, C.A. and Gerstein, A. 1997. Varying effects of low dissolved oxygen on trophic interactions in an estuarine food web. *Ecological Monographs*, 67(4), pp.489-507.

Diaz, R.J. and Rosenberg, R. 1995. Marine benthic hypoxia: a review of its ecological effects and the behavioral responses of benthic macrofauna. *Oceanography and marine biology. An annual review*, 33, pp.245-03.

Section VIII: Habitat

Hare JA, Morrison WE, Nelson MW, Stachura MM, Teeters EJ, et al. 2016. A Vulnerability Assessment of Fish and Invertebrates to Climate Change on the Northeast U.S. Continental Shelf. *PLoS ONE* 11(2): e0146756. doi: 10.1371/journal.pone.0146756.

Lee, M., E. Shevliakova, S. Malyshev, P.C.D. Milly, and P.R. Jaffé. 2016. Climate variability and extremes interacting with nitrogen storage, amplify eutrophication risk, *Geophys. Res. Lett.*, 43 7520-7528, doi:10.1002/2016GL069254.

Zimmerman, R.C., Hill, V.J. and Gallegos, C.L. 2015. Predicting effects of ocean warming, acidification, and water quality on Chesapeake region eelgrass. *Limnology and Oceanography*, 60(5), pp.1781-1804.

Jarvis, J.C., Brush, M.J., Moore, K.A. 2014. Modeling loss and recovery of *Zostera marina* beds in the Chesapeake Bay: The role of seedlings and seed-bank viability. *Aquatic Botany* 113, 32-45.

Vaquer-Sunyer, R., Duarte, C.M. 2011. Temperature effects on oxygen thresholds for hypoxia in marine benthic organisms. *Global Change Biology* 17, 1788-1797.

Codispoti, L.A. 2010. Interesting times for marine N₂O. *Science*, 327(5971), pp.1339-1340.

Najjar, R.G., Pyke, C.R., Adams, M.B., Breitburg, D., Hershner, C., Kemp, M., Howarth, R., Mulholland, M.R., Paolisso, M., Secor, D. and Sellner, K. 2010. Potential climate-change impacts on the Chesapeake Bay. *Estuarine, Coastal and Shelf Science*, 86(1), pp.1-20.

Orth, R.J., Williams, M.R., Marion, S., Wilcox, D.J., Carruthers, T.J.B., Moore, K.A., Kemp, W.M., Dennison, W.C., Rybicki, N., Bergstrom, P., Batiuk, R. 2010. Long-term trends in submersed aquatic vegetation (SAV) in Chesapeake Bay, USA, related to water quality. *Estuaries and Coasts* 33, 1144–1163.

Palmer MA, DP Lettenmaier, NL Poff, SL Postel, B Richter, and R Warner. 2009. Climate change and river ecosystems: protection and adaptation options. *Environmental Management* 44: 1053-1068.

Moore, K.A. and Jarvis, J.C. 2008. Environmental factors affecting recent summertime eelgrass diebacks in the lower Chesapeake Bay: implications for long-term persistence. *Journal of Coastal Research*, pp.135-147.

Pyke, C.R., Najjar, R.G., Adams, M.B., Breitburg, D., Kemp, M., Hershner, C., Howarth, R., Mulholland, M., Paolisso, M., Secor, D. and Sellner, K. 2008. *Climate Change and the Chesapeake Bay: State-of-the-Science Review and Recommendations. A Report from the Chesapeake Bay Program Science and Technical Advisory Committee (STAC)*. Annapolis, MD, 59.

U.S. Environmental Protection Agency. 2008. *Climate Change and Aquatic Invasive Species*, Washington, D.C., EPA/600/R-08/014.

Boesch, D.F., Coles, V.J., Kimmel, D.G. and Miller, W.D. 2007. Coastal dead zones and global climate change: ramifications of climate change for Chesapeake Bay hypoxia. *Regional impacts of climate change: four case studies in the United States*, pp.57-70.

Palacios, S.L. and Zimmerman, R.C., 2007. Response of eelgrass *Zostera marina* to CO₂ enrichment: possible impacts of climate change and potential for remediation of coastal habitats. *Marine Ecology-Progress Series*, 344, p.1.

Twilley, R.R., Kleypas, J.A., Cowan Jr, J.H. and Hare, S.R. 2002. *Coastal and marine ecosystems & Global climate change: Potential Effects on US Resources* (pp. 1-64). Arlington, VA: Pew Center on Global Climate Change.

Wood, R.J., Boesch, D.F. and Kennedy, V.S. 2002. Future consequences of climate change for the Chesapeake Bay ecosystem and its fisheries. In American Fisheries Society Symposium (Vol. 32, pp. 171-184).

Najjar, R.G., Walker, H.A., Anderson, P.J., Barron, E.J., Bord, R.J., Gibson, J.R., Kennedy, V.S., Knight, C.G., Megonigal, J.P., O'Connor, R.E. and Polsky, C.D. 2000. The potential impacts of climate change on the mid-Atlantic coastal region. *Climate Research*, 14(3), pp.219-233.

Breitburg, D.L., Loher, T., Pacey, C.A. and Gerstein, A. 1997. Varying effects of low dissolved oxygen on trophic interactions in an estuarine food web. *Ecological Monographs*, 67(4), pp.489-507.

Moore KA, HA Neckles, and RJ Orth. 1996. *Zostera marina* (eelgrass) growth and survival along a gradient of nutrients and turbidity in the lower Chesapeake Bay. *Marine Ecology Progress Series* 142: 247-259.

Diaz, R.J. and Rosenberg, R. 1995. Marine benthic hypoxia: a review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanography and marine biology. An annual review*, 33, pp.245-03.

Section IX: Tools

Chesapeake Atlantis Model

The Chesapeake Atlantis Model was developed to conduct a simulation that accounts for tradeoffs in performance across a range of management objectives by incorporating spatially explicit information about the biological, geochemical, and physical dynamics of the Chesapeake Bay. This model has strong applications for simulating the direct and indirect impacts of climate change on the Bay ecosystem.

North Atlantic Landscape Conservation Cooperative Fish Habitat Decision Support Tool

This tool was created with funding from the United States Fish and Wildlife Service to provide resource managers and the general public with access to the extensive spatial data and results produced from multiple fish habitat assessments.

National Fish Habitat Partnership Data System

The National Fish Habitat Partnership (NFHP) data system supports coordinated efforts of scientific assessment and data exchange among the partners and stakeholders of the aquatic habitat community. Under the guidance of the NFHP Science and Data Committee, the system provides data access and visualization tools for authoritative NFHP data products and contributed data from partners. The underlying data management tools and best practices are

the foundation enabling partners, researchers, and managers to access and use data now and in the future.

[LandScope](#)

LandScope Chesapeake fills a need identified for a publicly accessible, watershed-wide land conservation priority system. Its purpose is to support collaboration among many partners in land conservation efforts throughout the region.

[Maryland GreenPrint](#)

The GreenPrint map displays targeted ecological areas (TEAs)—lands and watersheds of high ecological value that have been identified as conservation priorities by the Maryland Department of Natural Resources. It also displays information about four of Maryland's most active state-operated land conservation programs: Program Open Space - Stateside, the Maryland Agricultural Land Preservation Foundation, the Maryland Environmental Trust, and the Rural Legacy Program. GreenPrint offers a way to improve how these programs work together and assists in steering Project Open Space acquisitions to the TEAs. The TEAs were developed in 2008 and updated in 2011. This map shows the 2011 version of the TEAs, so some older acquisitions may not appear in the TEAs, even though they were acquired using the TEAs available at that time. The maps also display other protected land data for reference.

[Virginia Coastal Geospatial and Educational Mapping System](#)

Virginia Coastal Geospatial and Educational Mapping System (GEMS) is a gateway to Virginia's coastal resource data and maps; coastal laws and policies; facts on coastal resource values; and direct links to collaborating agencies responsible for current data. It includes a growing inventory of water- and land-based natural resources, conservation planning tools, and planning examples that can be used to help to protect Virginia's coastal ecosystems. Additionally, this tool was developed to promote community involvement and environmental education.

[Virginia Department of Conservation and Recreation Conservation Lands Database](#)

The Virginia Department of Conservation and Recreation digitally maps the boundaries for agency owned and managed state parks and state natural area preserves, the Commonwealth's state-wide Conservation Lands Database, and has grown to include state, federal, private, and locally managed lands and conservation easements.

[Virginia Department of Environmental Quality Virginia Environmental Geographic Information System \(VEGIS\) Database](#)

This application displays cross-media geographical features including petroleum release sites, solid waste facilities, 303D impaired waters, water-quality monitoring stations and more.

[Virginia Department of Game and Inland Fisheries Database](#)

The Mapping and Geographic Information Systems (GIS) section provides cartographic services, GIS/GPS training and technical support, spatial modeling, analysis, and geographic data development. The GIS section is within the Bureau of Wildlife Resources. The section supports a wide range of Department of Game and Inland Fisheries core functions, contributing map products and services throughout the agency. In addition, the GIS staff maintains internet mapping services for the public and provide data to other governmental and nongovernmental organizations.

[Habitat Prioritization Tool](#)

The Chesapeake Bay Habitat Tool provides spatial context for multi-habitat conservation opportunities in the Bay. Features of the tool include: a web map with Bay-wide data for important nearshore habitats; new benthic data layers to better describe the habitat on the Bay floor; and an interactive tool to prioritize wetland protection and restoration. This map and prioritization tool are designed to be screening-level tools. Results are not intended to be a replacement for site-specific knowledge nor a prescription for on-the-ground action.

[Chesapeake Bay Fish Passage Prioritization Tool](#)

The Chesapeake Fish Passage Prioritization project was designed by The Nature Conservancy to help managers identify potential fish passage projects that are most likely to produce ecological benefits. Results include three consensus-based scenarios that depict dams in the 64,000 square mile Chesapeake Bay watershed where passage projects would provide the greatest potential benefit for diadromous fish, resident fish, and brook trout, respectively. Additionally, an interactive map and tool allow users to examine results in the context of other relevant data, develop custom scenarios, and to model the effects of conducting a passage project at a given dam.

[Chesapeake Bay U.S. Geological Survey Data](#)

The data page contains USGS information for key national data sets and also more specific information for the Chesapeake Bay watershed. Links to selected state information are also included.