

N Application on Full Season Soybean

Region	Acres	Lbs Inorganic N	Lbs Manure N	Lbs N/ac	% N from Manure
Central	22,744	201,998	10,912	9	5%
Lower Eastern Shore	86,999	148,548	297,119	5	67%
North Western	63,390	420,716	271,941	11	39%
Northern	29,890	189,521	20,960	7	10%
Southern	15,400	160,717	6,323	11	4%
Upper Eastern Shore	108,576	607,229	127,574	7	17%
Western	1,797	9,337	12,006	12	56%

Average Statewide Load 2.5 lbs manure N (42% of N applied from manure)

Average Eastern Shore Load 3.4 lbs manure N (66% of N applied from manure)

P Application on Full Season Soybean

Region	Acres	Lbs Inorganic P2O5	Lbs Manure P2O5	Lbs P2O5/ac	% P from Manure
Central	22,744	387,972	12,253	18	3%
Lower Eastern Shore	86,999	259,783	343,346	7	57%
North Western	63,390	1,112,391	305,137	22	22%
Northern	29,890	708,596	18,108	24	2%
Southern	15,400	350,777	3,608	23	1%
Upper Eastern Shore	108,576	1,342,380	166,721	14	11%
Western	1,797	32,352	12,896	25	28%

(34% of N applied from manure)

(57% of N applied from manure on Lower Shore)