



Lessons from Chesapeake Bay restoration efforts:

Understanding the role of nutrient reduction activities in improving water quality

Bill Dennison

Christina Lyerly

University of Maryland Center for Environmental Science

Scientific, Technical Assessment and Reporting Meeting

November 20, 2013

Authors: Christina M. Lyerly, Ana L. Hernández Cordero, Katherine L. Foreman and William C. Dennison


Lessons from Chesapeake Bay Restoration Efforts

- Seven lessons under three broad categories:
 - What Works**
 - Challenges**
 - What We Need**
- Presented at the Coastal and Estuarine Research Federation conference (November 5th)
- Accepted as a poster presentation at the December 2013 Maryland Water Monitoring Council conference
- Scheduled for release January 2014




Lesson 1

- Upgrades in both nitrogen and phosphorus wastewater treatment result in rapid local water quality improvements



Trends in Wastewater Treatment Plant Loads

- Wastewater treatment plants yet to be upgraded represent opportunities for continued improvement




Upper Patuxent River

- Upgrades to a wastewater treatment plant reduced nitrogen loads to the Upper Patuxent River

DRAFT

Changes in TN and
TP loads
(1984-2004)




Upper Patuxent River

- Decreased wastewater treatment plant nutrient loads contributed to a resurgence of submerged aquatic vegetation



Vallisneria americana (wild celery). Photo © Cassie Gurbisz




Data from Boynton et al., 2008

Upper Potomac River


- Submerged aquatic vegetation abundance increased
- Total nitrogen concentrations decreased at the Piscataway Creek monitoring station

Changes in total SAV abundance in the Upper Potomac River (1990-2008)



Data from Ruhl and Rybicki, 2010

Changes in mean monthly surface TN concentrations at a monitoring station near Piscataway Creek (1985-2013)




Data from Romano and Buchanan, 2011

Lesson 2


- Improvements in air quality lead to reductions in atmospheric nitrogen deposition

Annual mean wet inorganic nitrogen deposition



Point Source Emissions and Surface Water Quality

- Decreases in atmospheric point source emissions and nitrogen deposition are linked to increased surface water quality in 9 mostly forested subwatersheds




Changes in nitrate-N concentrations at 3 water quality monitoring stations (1986-2009)

Nonpoint Source Emissions

- Mobile sources represent the majority of NO_x air emissions

U.S. NO_x emissions by category (2006)



Pre 1970s. Over concern from early smog studies in Los Angeles, CA, the catalytic converter was invented in the 1950s by Eugene Houdry, a French mechanical engineer living in the United States. Catalytic converters were further developed by John J. Mooney and Carl D. Keith, creating the first production catalytic converter in 1973.



1975. The U.S. EPA requires that all new cars be equipped with catalytic converters. These two-way converters combine carbon monoxide (CO) and unburned hydrocarbons (HC) to produce carbon dioxide (CO_2) and water (H_2O).



1981. To keep up with the Clean Air Act of 1970, a new generation of catalytic converters act as a three-way catalyst, converting carbon dioxide (CO_2) and hydrocarbons (HC) to water (H_2O), and reducing nitrogen oxides (NO_x) to elemental nitrogen (N) and oxygen (O).

Lesson 3

- Reductions of agricultural nutrient sources result in improved stream quality

Cover crops



Livestock exclusion




Waste management




Wye River Drainage Basin

- Winter cover crops planted in the Wye River drainage basin improved water quality



Changes in groundwater nitrate-N concentrations in 2 agricultural fields (1986-1998)



Changes in nitrate-N concentrations in different subsurface zones on an agricultural field (1992-1999)

Practices for Further Study


- Rotational grazing may provide some benefits but has shown mixed results
- Cattle feed composition may influence the volatilization of ammonia from animal waste
- Ventilation construction may affect ammonia emitted from poultry houses



Photo © jlastras

Lesson 4

- Many practices provide initial water quality improvements in runoff; however, full benefits to stream conditions can be delayed




Delmarva Peninsula

- Groundwater ages can range from less than a year to more than 100 years


Simulated return time of groundwater traveling from the water table to its discharge location

DRAFT



Lesson 5


- Improvements in water quality can be counteracted by changes in nutrient sources and land-use practices



Choptank River

- Wastewater treatment plant flow has increased in some locations, but WWTP upgrades have decreased nutrient loads


Changes in flow, N and P loads
from the Easton wastewater
treatment plant
(1995-2012)



Choptank River

- Agricultural nutrient sources have counteracted reductions in wastewater treatment plant nutrient loads

Increases in TN and TP at
Greensboro water quality
monitoring station
(1968-2012)



Lesson 6


- Observable water quality responses are more likely to occur if A) location specific sources of pollution are identified and B) targeted practices are implemented.



Corsica River


- Improvements in nontidal water quality in the Corsica River were observed after aggressive implementation of multiple nutrient reduction practices

Changes in TN and TP concentrations in Three Bridges Branch and Gravel Run (2006-2011)




Anne Arundel County

Location matters



Tidal Boundary Streams: 2 out of 3 restored streams retained nitrogen...Partial success



Headwaters Streams: 0 out of 3 restored streams retained nitrogen...No success

Positive Export Values = No Nitrogen Retention = Restored Streams Not Doing Their Jobs!

Lesson 7

- An array of practices to promote stormwater infiltration and retention are needed in urban and suburban areas

Rain gardens



Gutters and downspouts installed onto buildings and in lawns help assist in directing rain water from the roof to the garden. A landscape of native, drought resistant plants is well adapted to local conditions and easily maintained. Plants with deep root systems encourage stormwater infiltration and help absorb excess nutrient runoff. Additionally, a berm on the downward slope of a rain garden will help hold water in the garden during heavy rains, further improving its filtering capacity.

Pervious surfaces




Impervious surfaces such as cement, asphalt and roofing prevent the infiltration of stormwater, increasing the volume and velocity of surface runoff which carries nutrients and sediments with it. Pervious surfaces, such as pervious pavement or pavers, allow for stormwater to filter through the surface and into the ground, rather than into nearby streams and storm drains.

Constructed Wetlands

- Constructed wetlands in Baltimore, MD demonstrated the potential to reduce nitrate entering streams through stormwater runoff

Mean groundwater nitrate-N concentrations (A) and denitrification rates (B) in oxbow, forested, and constructed wetlands in Baltimore, MD



Conclusions

1. What Works



- The Clean Water Act works
- The Clean Air Act works
- Multiple practices that reduce agricultural nutrient loads work

2. Challenges

- Delays in improvements necessitate patience, persistence and perspiration
- We are not trying hard enough

3. What We Need

- Location, location, location should guide restoration efforts
- Innovative practices are needed to up our game



Thank you

Questions?



References

- Batchelor, T., C. Decker, F. Digiliano, J. Dindinger, Q. Forrest et al. 2011. Corsica River Targeted Initiative Progress Report: 2005-2011. Corsica River Initiative, 21 p.
- Boynton, W.R., J.D. Hagy, J.C. Cornwell, W.M. Kemp, S.M. Greene, M.S. Owens, J.E. Baker, and R.K. Larsen. 2008. Nutrient budgets and management actions in the Patuxent River Estuary, Maryland. *Estuaries and Coasts* 31: 623-651.
- Eshleman, K.N., R.D. Sabo, and K.M. Kline. 2013. Surface water quality is improving due to declining atmospheric N deposition. *Environmental Science & Technology*.
- Filoso, S. and M.A. Palmer. 2009. Case Study #4: Assessing restoration effectiveness at reducing nitrogen loads. In 2010 Trust Fund: Water Quality Monitoring Strategy, Section 4, 4 p.
- Fisher, T.R., J.D. Hagy III, W.R. Boynton, and M.R. Williams. 2006. Culture eutrophication in the Choptank and Patuxent estuaries of Chesapeake Bay. *Limnology and Oceanography*, 51: 435-444.
- Harrison, M.D., P.M. Groffman, P.M. Mayer, S. Srinivasan, and T. J. Corns. 2011. Denitrification in alluvial wetlands in an urban landscape. *Journal of Environmental Quality* 40: 640-646.
- Karrh, R. 2012. Choptank, Little Choptank and Patuxent Rivers Water Quality and Habitat Assessment. *Maryland Department of Natural Resources Tidewater Ecosystem Assessment*, 6 p. Available at: <http://mddnr.chesapeakebay.net/eyeson/ebay/documents/Choptank%20Summary%20WQ%20and%20H%20Assessment%202012.pdf>.
- Phillips, S.W. and B.D. Lindsey. 2003. The influence of ground water on nitrogen delivery to the Chesapeake Bay. US Geological Survey Fact Sheet FS-091-03, 6 p.
- Sanford, W.E., J.P. Pope, D.L. Selim, and R.F. Stumvoll. 2012. Simulation of groundwater flow in the shallow aquifer system of the Delmarva Peninsula, Maryland and Delaware. U.S. Geological Survey Open-File Report 2012-1140, 58 p. Available at: http://pubs.usgs.gov/of/2012/1140/pdf/OFR_2012-1140.pdf.
- Staver, K. and R. Brinsfield. 1995. Case Study #1: Assessing the impact of changes in management practices on nutrient transport from coastal plain agricultural systems. In 2010 Trust Fund: Water Quality Monitoring Strategy. 3 p. Available at: http://www.dnr.state.md.us/ccp/funding/pdfs/Monitoring_Strategy.pdf.
- Staver, K. and R. Brinsfield. 2000. Evaluating changes in subsurface nitrogen discharge from an agricultural watershed into Chesapeake Bay after implementation of a groundwater protection strategy. In 2010 Trust Fund: Water Quality Monitoring Strategy. 3 p. Available at: http://www.dnr.state.md.us/ccp/funding/pdfs/Monitoring_Strategy.pdf.
- Testa, J.M., W.M. Kemp, W.R. Boynton, and J.D. Hagy III. 2008. Long-term changes in water quality and productivity in the Patuxent River Estuary: 1985 to 2003. *Estuaries and Coasts* 31: 1021-1037.
- U.S. Environmental Protection Agency Office of Transportation and Air Quality. 2008. 2007 Progress Report: Vehicle and Engine Compliance Activities, EPA-420-R-08-011, 59 p.
- U.S. Environmental Protection Agency. 2009. Clean Air Markets 2009 Environmental Results. Available at: http://www.epa.gov/airmarkt/progress/ARP09_3.html. Accessed on April 25, 2013.