


# **Estimating Floodplain Benefits in the Chesapeake**

Krissy Hopkins, Ph.D. U.S. Geological Survey

Scientific, Technical Assessment and Reporting (STAR) Meeting
May 23, 2024



# **Study Objective**

What is the capacity of floodplains to **retain sediment**, **nutrients**, and flood waters?






# **Study Area**

# Floodplain Sediment and Nutrient Retention

Non-tidal wadable streams in the Chesapeake and Delaware River watersheds.






#### **Methods Workflow**

# Field Geomorphic Measurements


Floodplain deposition and streambank erosion



#### Watershed + Reach Predictors


#### Reach fluvial geomorphometry

Floodplain width, bank height, channel width Hopkins et al. 2020





# Floodplain Deposition Measurements



Site Layout: 100 m reach



#### Field survey metrics

Active floodplain width

Bank height

Channel width

Lateral and vertical change (cm/yr)


% eroding bank

Adjusted lateral erosion (cm/yr)

#### Floodplain deposition




562 floodplain tree cores with mean tree age of 48 years.



Noe et al. 2020

### Streambank Erosion Measurements



Site Layout: 100 m reach



#### Field survey metrics

Active floodplain width

Bank height

Channel width

Lateral and vertical change (cm/yr)


% eroding bank

Adjusted lateral erosion (cm/yr)

#### **Streambank Erosion**



492 root cookies with mean exposure age 17 years.




Noe et al. 2020

https://doi.org/10.5066/P93OUWYZ

## Sediment Characteristics at 68 Sites

296 Floodplain samples = 461 Stream bank samples =







Bulk density (g/cm<sup>3</sup>)

Bulk density <2 mm (g/cm<sup>3</sup>)

Bulk density <1 mm (g/cm<sup>3</sup>)

% organic

% mineral

% carbonate

Total OC (%)

Total N (%)

Total P (mg/g)

Total Ca (mg/g)

Total Na (mg/g)

Total Mg (mg/g)

Total K (mg/g)

Total AI (mg/g)

Total Fe (mg/g)

Total Ti (mg/g)

Particle size: mean (um)

Particle size: d50 (um)


Particle size: %<63 um



## **Sediment Flux**

Floodplain Flux = vertical accretion rate (m/yr) x floodplain width (m) x sediment bulk density  $(kg/m^3) = kg/m/yr$ 

Bank Flux = lateral erosion rate x bank height x 2 banks x sediment bulk density






### **Methods Workflow**

# Field Geomorphic Measurements


Floodplain deposition and streambank erosion



#### Watershed + Reach Predictors

#### Reach fluvial geomorphometry

Floodplain width, bank height, channel width Hopkins et al. 2020





# Floodplain and Chanel Evaluation Tool (FACET)


Automated fine-scale fluvial geomorphometry applied to large scales.

#### **Functionality**

- DEM pre-processing
- Stream network delineation
- Stream bank detection
- Mapping floodplain extent
- Calculate channel and floodplain metrics

#### **Metrics**


- Bank height
- Channel width
- Channel area
- Floodplain width








## **FACET Channel and Floodplain Dimensions**



### **Methods Workflow**

# Field Geomorphic Measurements


Floodplain deposition and streambank erosion



#### Watershed + Reach Predictors


#### Reach fluvial geomorphometry


Floodplain width, bank height, channel width Hopkins et al. 2020





### **Random Forest Models**









### Floodplain Deposition and Streambank Erosion Predictions

#### **Streambank Nitrogen Flux**



#### Floodplain Nitrogen Flux



Summarize conditions at river basin and catchment scales.

Identify areas more prone to erosion or deposition.


Useful in **targeting** areas for **restoration** or high-value areas for **conservation**.



### **Methods Workflow**

# Field Geomorphic Measurements


Floodplain deposition and streambank erosion



#### Watershed + Reach Predictors

#### Reach fluvial geomorphometry

Floodplain width, bank height, channel width Hopkins et al. 2020






## **Benefits Transfer Approach for Valuation**

Treatment costs to reduce nitrogen from 4.0 to 7.5 mg/L **\$16.12 USD per kg-N.** 



Blue plains wastewater treatment plant.

# Study within the Chesapeake Bay watershed in Virginia.





Hopkins et al. 2018

https://doi.org/10.1016/j.jenvman.2018.05.013

Hopkins et al. 2023

https://doi.org/10.1016/j.jenvman.2023.118747

# Floodplains Provide Substantial Benefits by Trapping Sediments and Nutrients






Net
Nitrogen
Retention
\$100 million
USD per year

Chesapeake Bay watershed



# Spatial Variability in the Value of Floodplain Sediment and Nutrient Retention






# Coastal Plain Floodplains Provided the Highest Nutrient Trapping Benefits
















Conestoga River

# Headwater Streams Net Source of Sediment, but a Net Sink for Nitrogen











More nitrogen is deposited in headwater floodplains.



More sediment is eroded from streambanks than deposited on the floodplain.

# Higher Order Streams Provide the Most Nitrogen Trapping per Length of Stream







# Floodplain Ecosystem Services Mapper





# **Summary**

Floodplains provide substantial benefits by trapping nutrients and sediment, upwards of \$220 million per year in the Chesapeake Bay watershed.

