## Office Hours AMT

3/14/2025

#### Main topics

- Crops in other hay
- Poultry data (note this is being presented by DE)
- Inorganic Ag Fertilizer

#### February Recap: 5 Decisions

- 1. Two new Land Uses Managed Hay and Pasture.
- 2. Four manure application groups
- 3. Defined managed and unmanaged hay and pasture.
- 4. Acres of managed Land Uses will be state reporting each year by November 1<sup>st</sup>.
- 5. Assigned Non Nutrient Management multipliers for Nitrogen and Phosphorus.

# A quick caveat:

| Crop:  | CAST Land Use: |
|--|----------------|
| bromegrass seed                                      | Other Hay      |
| cropland on which all crops failed or were abandoned | Other Hay      |
| fescue seed  | Other Hay      |
| orchardgrass seed                                    | Other Hay      |
| other field and grass seed crops                     | Other Hay      |
| other haylage; grass silage and greenchop            | Other Hay      |
| other managed hay                                    | Other Hay      |
| ryegrass seed  | Other Hay      |
| small grain hay                                      | Other Hay      |
| timothy seed   | Other Hay      |

# A quick caveat:

#### Fixes Nitrogen

| Crop Name:   | CAST Land Use: |
|--|----------------|
| bromegrass seed                                      | Cther Hay      |
| cropland on which all crops failed or were abandoned | Other Hay      |
| fescue seed  | Other Hay      |
| orchardgrass seed                                    | Other Hay      |
| other field and grass seed crops                     | Other Hay      |
| other haylage; grass silage and greenchop            | Other Hay      |
| other managed hay                                    | Other Hay      |
| ryegrass seed  | Other Hay      |
| small grain hay                                      | Other Hay      |
| timothy seed   | Other Hay      |

#### Some things to think about


- Other haylage; grass silage and greenchop
  - Ag census definition:
    - "haylage, grass silage, and greenchop (hay cut and fed green) made from such crops or mixtures as small grains, clover, lespedeza, fescue, timothy, Bermuda grass, sudangrass, sorghum-sudan crosses, orchardgrass, soybeans, peanuts, etc".
  - Currently:
    - 30.24 lbs N fixed per yield unit
    - Currently ~ 25% of other hay acres

#### Should we switch the Land Use?

| Crop Name:  | CAST Land Use: |
|---|----------------|
| alfalfa seed  | Leguminous Hay |
| vetch seed  | Leguminous Hay |
| alfalfa hay   | Leguminous Hay |
| birdsfoot trefoil seed                                | Leguminous Hay |
| haylage or greenchop from alfalfa or alfalfa mixtures | Leguminous Hay |
| red clover seed                                       | Leguminous Hay |
| other haylage; grass silage and greenchop             | Other Hay      |
| cropland on which all crops failed or were abandoned  | Other Hay      |
| other managed hay                                     | Other Hay      |
| orchardgrass seed                                     | Other Hay      |
| other field and grass seed crops                      | Other Hay      |
| ryegrass seed   | Other Hay      |
| fescue seed   | Other Hay      |
| bromegrass seed                                       | Other Hay      |
| timothy seed  | Other Hay      |
| small grain hay                                       | Other Hay      |

#### What is the impact of changing this?

Watershed Percentage of Other Hay acres that are OtherHaylageGrassSilageGreenchop



#### What is the impact of changing this?

| Year | Sum of OtherHayAcres | Sum of OtherHaylageGrassSilageGreenchopAcres |
|------|----------------------|--|
| 2002 | 2,365,400            | 236,489                                      |
| 2003 | 2,333,419            | 239,520                                      |
| 2004 | 2,301,479            | 242,551                                      |
| 2005 | 2,269,541            | 245,582                                      |
| 2006 | 2,237,605            | 248,613                                      |
| 2007 | 2,205,682            | 251,645                                      |
| 2008 | 2,189,357            | 262,781                                      |
| 2009 | 2,173,031            | 273,918                                      |
| 2010 | 2,156,706            | 285,054                                      |
| 2011 | 2,140,380            | 296,190                                      |
| 2012 | 2,124,544            | 307,696                                      |
| 2013 | 2,143,608            | 350,402                                      |
| 2014 | 2,164,158            | 393,389                                      |
| 2015 | 2,184,774            | 436,380                                      |
| 2016 | 2,205,391            | 479,371                                      |
| 2017 | 2,227,549            | 522,833                                      |

### Questions?

#### The discussion: Where do we want to go?

- Crop: other haylage; grass silage and greenchop
  - Current Land Use: Other Hay
  - Suggested Land Use: Leguminous Hay

#### Calculating manure nutrients



#### Poultry Industry Input

Animal population (AU)

Dry pounds of manure nutrient conc.

Manure nutrients

- Weights
- -Numbers


- Manure/litter generation

- Nutrient content

#### Recap inorganic fertilizer in CAST

Current Foundational Path towards fertilizer knowledge improvement scale

#### Foundational:




- CURRENTLY fertilizer from ALL states feeds a single stock
- Fertilizer from one state can theoretically be applied in another
- Implications for all of the watershed


### Basics of nutrient application




#### Basics of nutrient application



#### Basics of nutrient application



County A organic nutrient stock (biosolids, manure)





Percentage of crop A nutrient need met:


Crop A

Crop A


0%









Percentage of crop A nutrient need met:




Fertilizer stock for the entire Chesapeake Bay watershed



Percentage of crop need met:




Fertilizer stock for the entire Chesapeake Bay watershed



- This calculation can only be done when we have actual fertilizer data
- Provides an application ratio of crop need (organic nutrients to inorganic nutrients (50/50))
- This ratio is used to apply nutrients for all subsequent years

Varying yields impact applications



Percentage of crop A nutrient need met:

50%
inorgani
c
Crop A


100%

50%

organic


# Applications and crop yield data example: 100% of crop need is met


Contribution of organic and inorganic nutrients to Crop A as a function of yields over time





#### Some caveats:

- BMPs can impact application
  - Load Source Reduction (Core NM)
- BMPs can shift/replace nutrients
  - Manure transport
  - Manure treatment technologies











#### Remember, fertilizer is a watershed product

• If one area uses less fertilizer other areas will use more.



#### A note on Nitrogen Loading

If you have two IDENTICAL fields, and apply the same amount Inorganic fertilizer will load MORE than manure




Runoff from Fertilizer



**Runoff from Manure** 



#### Sensitivities – all else being equal



### Questions?

#### What can we improve?

#### Different scale?

States won't influence each other the same way.

### Different methods for backfilling?

 Cap the percentage of fertilizer used to replace organic nutrients.

# Thank you for attending office hours!

We will begin our main meeting at 09:00.