
Review of WQ Assessment with the 4-D interpolator

Gary Shenk
Bay Oxygen Research Group
9/9/2022

Motivation


92 segments




Up to 11 designated uses per segment

Refined Designated Uses for the Bay and Tidal Tributary Waters


A. Cross Section of Chesapeake Bay or Tidal Tributary



B. Oblique View of the "Chesapeake Bay" and its Tidal Tributaries



838 combinations




326 assessed




Zero segments fully assessed


The Plan – Create a 3D representation of water quality through time (4D)




WQ Assessment with 4D interpolator – big picture



WQ Assessment with 4D interpolator – big picture





WQ Assessment with 4D interpolator – big picture



Observed Data – Kitchen sink approach

Observed data





- Temperature
- Salinity
- Density
- ...

- Inflows
- Tide


Wind


 Modeled Hydrodynamics

•

Presentations today from Peter Tango and Jon Harcum

Mean Daily Prediction







- GAM = Generalized Additive Model
 - Rebecca Murphy and Elgin Perry presentations
- Produces the most likely water quality values for all space and time.
 - Dissolved oxygen
 - Clarity
 - Chlorophyll a
 - Pycnocline upper and lower boundary
 - Density => salinity and temperature

Presentation today from Elgin Perry

Extras

Statistical Simulation






- Statistical simulation
 - Deterministic addition of diel cycling
 - Stochastic autoregressive component to generate multiple feasible realizations
- incorporates observed spatial and temporal correlation.
- Data
 - vertical profiler
 - data flow
 - continuous monitoring data

WQ assessment – Why multiple realizations?

- Water Quality assessment is an estimation of the frequency of low-oxygen events
- Using the mean daily prediction would underestimate the frequency of these events
- The assessment is performed by calculating a frequency distribution of spatial violation rates
- The more available estimates of spatial violation rates, the more accurate the frequency distribution



tool

data

product


30-day mean – CB3MH – Deep Water

Table 2 Dissolved oxygen criteria and applicable designated uses of the water quality standards for Chesapeake Bay and its tidal tributaries. Assessed criteria and designated uses are set in italics and unassessed criteria and designated uses are in bold.

Designated use	Dissolved oxygen criteria concentration/duration		Temporal application
Migratory fish spawning and nursery use	7-day mean ≥ 6 m Instantaneous min	ng/L for tidal habitats with 0–0.5 ppt salinity nimum≥ 5 mg/L	February 1–May 31
	Open water fish and shellfish designated use criteria apply		June 1-January 31
Shallow water Bay grass use	Open water fish and shellfish designated use criteria apply		Year-round
Open-water fish and shellfish use	30-day mean	\geq 5.5 mg/L, salinity: 0–0.5 ppt \geq 5 mg/L, salinity: $>$ 0.5 ppt	Year-round
	7-day mean	$\geq 4 \text{ mg/L}$	
	Instantaneous min	$ \frac{1}{1} \frac{1} \frac$	
Deep-water seasonal fish and skellfish us	30 day mean > 3 mg		June 1–September 30
	Instantaneous minimum≥1.7 mg/L		
	Open-water fish and shellfish designated use criteria apply		October 1-May 31
Deep-channel seasonal refuge use	$Instantaneous\ minimum > 1\ mg/L$		June 1–September 30
	Open-water fish a	nd shellfish use criteria apply	October 1-May 31


Generate a spatially-complete realization of CB3MH-Deep Water oxygen 30-day mean concentration

T = day1-day30




Calculate the fraction of the volume below 3 mg/l

Produce an estimate for each summer month in 3 years




Typical 30-day mean curve: 4 months x 3 years




However, GAM results of the most likely value will underestimate the fraction of volume with extreme values

T = day1-day30



Need to have an accurate estimate of the fraction of the volume below 3 mg/l given the observations.

Enough realizations need to be produced to produce an accurate temporal distribution



A stochastic simulation could generate a more robust curve

