

High-resolution Landscape Characterization to Inform the Next Generation of Hydrologic Models

Peter R. Claggett¹, Labeeb Ahmed², Matthew Baker³, Jacob Czawlytco⁴, Sarah McDonald¹, Sean MacFaden⁵, Jarlath O'Neil-Dunne⁵, David Saavedra⁴, and Rachel Soobitsky⁴

1 Presenting Author, Lower Mississippi-Gulf Water Science Center, U.S. Geological Survey, Annapolis, MD 21403

2 Attain LLC, Annapolis, MD 21403

3 Department of Geography and Environmental Systems, University of Maryland, Baltimore County, MD 21250

4 Chesapeake Conservancy, Annapolis, MD 21403

5 University of Vermont Spatial Analysis Laboratory, Burlington, VT 05405

U.S. Department of the Interior U.S. Geological Survey




Chesapeake Bay 1-Meter Resolution Data

- 1. Land cover (12 classes): 2013/14, 2017/18, 2021/22
- 2. Land use (58 classes): 2013/14, 2017/18, 2021/22
- 3. Streams/ditches: best available LiDAR data
- 4. Stream channel and floodplain attributes: best available LiDAR data



Additional LiDAR Derivatives


- Height (normalized Digital Surface Model (nDSM), 1m LiDAR)
- Compound Topographic Index (3-meter, LiDAR)
- Topographic Wetness Index (1-meter, LiDAR)
- Multiscale Landforms (1-meter, LiDAR)











Local land use and parcel data

- Low-density Residential
- Recreation
- Agriculture
- Roads


High-resolution land cover data

- Impervious surfaces
- Tree canopy
- Low vegetation
- Water


CBP Land Uses

- Impervious-Roads
- Forests
- Turf Grass
- Open Space














2013/14 Chesapeake Bay Watershed Land Cover/Use Data

1m Land Cover



1m Land Use



10m Land Cover/Use


Impervious Roads Impervious Non-Roads Tree Canopy Over Impervious Tree Canopy Over Turf Grass **Turf Grass Forest** Wetlands (Floodplain) Wetlands (Other) Wetlands (Tidal) Water Mixed Open Cropland Pasture

https://chesapeake.usgs.gov/ph ase6/map/#map=7/-8582732.74/4851421.17/0.0/0,4



Chesapeake Bay Program Land Use Classification

1. Water (8)	2.2 Bare Construction	4.3 Extractive
1.1 Lentic	2.3 Turf Grass	4.3.1 Barren
1.1.1 Estuary	2.4 Tree Canopy over Turf Grass	4.3.2 Herbaceous
1.1.2 Lakes & Ponds		4.3.3 Scrub-shrub
	. Forest (5)	4.4 Solar fields
1.2.1 Streams	3.1 Contiguous (> 1 acre)	
1.2.1.1 Daylighted	3.2 Fragmented (< 1 acre)	5. Wetland (19)
1.2.1.2 Shaded	3.3 Natural Succession (e.g., Fallow)	5.1 Tidal
1.2.1.3 Culverted/ Buried	3.3.1 Barren	5.1.1 Open water
1.2.2.Ditches	3.3.2 Herbaceous	5.1.2 Barren
1.2.2.1 Daylighted	3.3.3 Scrub-shrub	5.1.3 Herbaceous
1.2.2.2 Shaded		5.1.4 Scrub-shrub
1.2.2.3 Culverted/ Buried 4	. Production (14)	5.1.5 Contiguous Forest
	4.1 Agriculture	5.1.6 Fragmented Forest
2. Developed (13)	4.1.1 Cropland	5.2 Non-tidal
2.1 Infrastructure	4.1.1.1 Barren	5.2.1 Floodplain/ Headwater
2.1.1 Roads	4.1.1.2 Herbaceous	5.2.1.1 Open water
2.1.2 Tree Canopy (TC) over Roads	4.1.2 Pasture	5.2.1.2 Barren
2.1.3 Structures	4.1.2.1 Barren	5.2.1.3 Herbaceous
2.1.4 TC over Structures	4.1.2.2 Herbaceous	5.2.1.4 Scrub-shrub
2.1.5 Other Impervious	4.1.3 Orchard/vineyard	5.2.1.5 Contiguous Forest
2.1.5.1 Railroads	4.1.3.1 Barren	5.2.1.6 Fragmented Forest
2.1.5.2 Parking Lots, Driveways, etc.	4.1.3.2 Herbaceous	5.2.2 Other
2.1.6 TC over Other Impervious	4.1.3.3 Scrub-shrub	5.2.2.1 Open water
2.1.6.1 TC over Railroads	4.2 Timber Harvest	5.2.2.2 Barren
2.1.6.2 TC over Parking Lots, etc.	4.2.1 Barren	5.2.2.3 Herbaceous
2.1.7 Suspended Succession	4.2.2 Herbaceous	5.2.2.4 Scrub-shrub
2.1.7.1 Barren	4.2.3 Scrub-shrub	5.2.2.5 Contiguous Forest
2.1.7.2 Herbaceous		5.2.2.6 Fragmented Forest
2.1.7.3 Scrub-shrub		5.3 Bare shore
≥USGS		575 Bail 5 51161 5



Tree Canopy Change in Two Suburban Counties

Prince George's County: 2014 - 2018

TC Loss (7,673 acres):

- 59% of loss change occurred within forest or wetlands
- 41% of loss occurred in developed areas

TC Gain (518 acres):

- 16% of gain occurred within forest or wetlands
 - shrub/scrub; edge of forest
- 54% of gain occurred in developed areas
- 29% of gain occurred on agricultural lands

Anne Arundel County: 2014 - 2018

TC Loss (2,544 acres):

- 57% of loss change occurred within forest or wetlands
- 42% of loss occurred in developed areas

TC Gain (188 acres):


- 9% of gain occurred within forest or wetlands
 - shrub/scrub; edge of forest
- 55% of gain occurred in developed areas
- 35% of gain occurred on agricultural lands




2017 Land Cover Production Status G cpo ford warranty - Google Searc ★ cicqis.org/portal/apps/webappviewer/index.html?id=262ce838a60048e9a0f136d904639f66 Land Cover Production Schedule and Hydrography Status with Web AppBuilder for ArcGIS Nashua Find address or place Albany Boston Layer List Worcester Springfield α≅ Layers Plymouth \bigcirc Providence Hartford ▼ Hydrography Status ••• Waterbury Cleveland Hydrography ... Bridgeport Island Completed (Subject to revisions) Youngstown Completed (Automated only, no QAQC) New York Queued Edison Not processed Pittsburgh Trenton Draft Land Cover Production Schedule Philadelphia Toms River Chesapeake Bay Watershed Atlantic City Land Cover Production Schedule ... April 2020 May 2020 June 2020 July 2020 Charleston August 2020 September 2020 October 2020 November 2020 -78.245 37.173 Degre of Chester, Esri, HERE, Garmin, FAO, NOAA, USGS, EPA, NPS | County of Ches...

Search for anything

USGS Land Change Monitoring, Assessment, and Projection Data Thirty Years of Change (1985 – 2015)











- 1. Lidar elevation
- 2. Valley-scale geomorphons
- 3. Channel-scale geomorphons
- 4. Extract valley network
- 5. Extract channels using valley network
- 6. QAQC channel skeleton
- 7. Connect stream network



^{* 1-}meter raster, 1:2000 scale

Hyper-Res Hydrography Production cicqis.org/portal/apps/webappviewer/index.html?id=262ce838a60048e9a0f136d904639f66 Land Cover Production Schedule and Hydrography Status with Web AppBuilder for ArcGIS Nashua Find address or place Albany Boston Layer List Worcester Springfield Q≅ Layers \bigcirc Providence Hartford Hydrography Status ••• Waterbury Cleveland ▼ ✓ Hydrography ••• Bridgeport Island Completed (Subject to revisions) Youngstown Completed (Automated only, no QAQC) New York Queued Allentown Edison Not processed Pittsburgh Trenton ▼ Draft Land Cover Production Schedule Philadelphia Toms River ▼ Chesapeake Bay Watershed ... Atlantic City Land Cover Production Schedule April 2020 May 2020 June 2020 July 2020 Charleston August 2020 September 2020 October 2020 November 2020 -82.277 41.611 De of Chester, Esri, HERE, Garmin, FAO, NOAA, USGS, EPA, NPS | Chesapeake C... All rights reserved Search for anything


USGS Floodplain and Channel Evaluation Toolkit




Channel Reach Measures

- Bank height (m)
- Bank angle, avg (deg)
- Bank angle, max (deg)
- Channel width (m)
- Channel length (m)
- Bankfull area (m²)
- Sinuosisty
- Drainage area (km²)
- Slope (%)
- Floodplain width (m)
- Floodplain elevation, range (m)
- Floodplain elevation, sd (m)





























Chesapeake Bay Land Change Model v5

Vulnerable farmland:
Area surrounded by recent residential and commercial growth



Summary

Under a 6-year Cooperative Agreement with the Chesapeake Bay Program, the Chesapeake Conservancy (CC) and University of Vermont (UVM) are producing 1-meter resolution land cover datasets for the years 2013/14, 2017/18, 2021/22 using the best available LiDAR and NAIP aerial imagery for the Bay watershed (including all of Maryland).

The CC and UVM are translating 12-classes of land cover into 58 of land use using ~15 ancillary datasets for each county.

The CC and UMBC are developing 1-meter resolution stream and ditch data, complementing the 1-meter resolution water bodies identified in the land cover data.

USGS has developed a GIS tool (FACET) to extract channel width, height, bank angle, and floodplain width and volumes from LiDAR.



Chesapeake Bay 1-Meter Resolution Release Dates

Land cover (12 classes, 1-meter):

- 2013/14 (available now- https://chesapeakeconservancy.org/conservation-innovation-center/high-resolution-data/land-cover-data-project/)
- 2017/18 (summer 2021)
- 2021/22 (summer 2024)

Land use (16 classes, 1-meter):

2013/14 (available now- https://chesapeakeconservancy.org/conservation-innovation-center/high-resolution-data/land-use-data-project/)

Land use (13 classes, 10-meter):

2013/14 (available now- https://chesapeake.usgs.gov/phase6/map/#map=7/-8582732.74/4851421.17/0.0/0,4)

Land use (58 classes):

- 2013/14 (summer 2021)
- 2017/18 (summer 2021)
- 2021/22 (summer 2024)

Streams/ditches

- Draft (summer 2021)
- Final (summer 2023)

Stream channel and floodplain attributes (FACET)

- Version 1 (available now- https://www.sciencebase.gov/catalog/item/5cae39c3e4b0c3b00654cf57
- Version 2 (2022 2023)



