Annual Phosphorus Loss Estimator (APLE) Model

Guido Yactayo - UMCES


APLE Model Capabilities

| | APLE |
|--|-------------------------------------|
| Language | It is a Microsoft Excel spreadsheet |
| Time Step | Annual |
| Units | Metric/ US |
| Edge-of-field P loss | YES |
| Simulates dissolved phosphorus loss in surface runoff | YES |
| Simulates sediment bound phosphorus loss in surface runoff | YES |
| Consider subsurface loss of P through leaching or artificial drainage. | NO |
| Consider all kinds of animal manure, applied either by machine or by grazing cattle, but consider only highly soluble commercial fertilizers | YES |
| Simulate P loss through grassed waterways or buffers that may occur beyond the field edge | NO |

APLE Model Inputs

| | APLE |
|--|-----------------------|
| Texture,organic matter and soil bulk density | YES |
| The type of soil cover | NO |
| The initial concentration of soil Labile P | YES |
| Day of manure or fertilizer applications | NO |
| Manure applications | Wet manure |
| For fertilizer applications | P2O5 applied per acre |
| Precipitation and Runoff | Annual |
| Temperature | - |
| Sediment | Annual |
| When grazing animals are present | YES |

Diagram of APLE Nutrient Inputs and Soil Pools



Soil test P Datasets

 AgriAnalysis Lab (2003-2014) for all zip codes in the Chesapeake Bay


 Spectrum Analysis (2006-2013) data are at the scale of state

 Penn State (2001-2014) data is comparable to the AgriAnalysis data


Lancaster PA – high till with manure (hwm)

| YEAR | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| ANNUAL_RAIN_IN | 37.7 | 48.8 | 49.0 | 41.5 | 60.1 | 33.0 | 43.2 | 45.5 | 42.3 | 32.0 | 44.2 | 58.1 | 54.5 | 46.6 |
| ANNUAL_RUNOFF_IN | 0.2 | 2.2 | 0.4 | 0.4 | 4.2 | 0.6 | 0.4 | 1.6 | 1.5 | 0.4 | 0.2 | 3.8 | 1.9 | 2.1 |
| SEDIMENT TONACRE | 0.9 | 21.6 | 1.5 | 1.7 | 30.6 | 3.2 | 2.4 | 25.8 | 13.2 | 1.5 | 0.5 | 10.3 | 19.5 | 34.6 |

P Loss in Runoff



Mehlich 3 Soil P




Somerset MD – high till with manure (hwm)

| YEAR | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 |
|------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| ANNUAL_RAIN_IN | 47.0 | 46.0 | 46.7 | 37.1 | 57.5 | 41.6 | 44.6 | 50.2 | 50.8 | 46.1 | 48.7 | 62.0 | 51.6 | 49.3 |
| ANNUAL_RUNOFF_IN | 2.5 | 3.8 | 4.3 | 2.2 | 6.2 | 2.2 | 4.2 | 5.6 | 6.3 | 5.8 | 4.7 | 7.8 | 6.6 | 6.4 |
| SEDIMENT_TONACRE | 1.3 | 1.7 | 2.4 | 1.7 | 2.7 | 1.1 | 2.1 | 2.4 | 3.8 | 2.9 | 2.6 | 4.1 | 2.4 | 3.8 |

P Loss in Runoff




Mehlich 3 Soil P




APLE Sensitivity to fertilizer inputs – Somerset MD (hwm)


Total P loss




Final Mehlich 3 Soil P Entire Soil



APLE sensitivity to fertilizer inputs - Lancaster PA (hwm)





APLE Model in Phase 6

- APLE has been Implemented in Phase 6
- Soil P test data has been acquired
- Long term sensitivity analysis vs a 10 year analysis?