

NOAA's Sea Level Rise Viewer

And Marsh Resilience

Nate Herold nate.herold@noaa.gov


Outline


- Introduction to NOAA's Sea Level Rise Viewer
 - Overview of the viewer
 - Marsh Migration (and impact) modeling
- Marsh Resilience Analysis
 - Landscape Scale / national assessment
 - Local level assessments
- Coastal Change Analysis Program (C-CAP) Land Cover
 - High Resolution future



NOAA's Sea Level Rise Viewer











Sea Level Rise Viewer



Sea Level Rise


View sea level rise and potential coastal flooding impact areas and relative depth.


• 0 0 0 0 0


GET STARTED


Disclaimer


The data and maps in this tool illustrate the scale of potential flooding, not the exact location, and do not account for erosion, subsidence, or future construction. Water levels are relative to Mean Higher High Water (MHHW) (excludes wind driven tides). The data, maps, and information provided should be used only as a screening-level tool for management decisions. As with all remotely sensed data, all features should be verified with a site visit. The data and maps in this tool are provided "as is," without warranty to their performance, merchantable state, or fitness for any particular purpose. The entire risk associated with the results and performance of these data is assumed by the user. This tool should be used strictly as a classified results and performance of the second of the provided results are relative to Mean Higher High Water (MHHW) (excludes wind driven tides).




























































































Direct Comparison of Data Layers

2016 Land Cover

2060 Intermediate High


with high rate of accretion (1ft Net Impact)





Marsh Resilience Analysis








National Marsh Resilience Metrics

	Group	Metric	+/-	Explanation
	Marsh Configuration	Area to edge ratio	+	more core and less edge represents less exposure
Current		Unvegetated edge ratio - more unvegetated edge represents more risk	more unvegetated edge represents more risk	
Condition		% impervious	-	more impervious can negatively impact runoff/habitat
Condition	Land Cover/Use	% natural	+	more natural areas can absorb runoff/protect habitat
		% agricultural	-	more agriculture can negatively impact runoff/habitat
	Erosion	Soil erodibility	-	high erosion potential represents greater risk of loss
Vulnerability		Tidal range	+	a larger tidal range represents more resilience
Valifierability	Tides	% marsh below MHHW	-	more marsh below MHHW represents more risk with SLR
		% marsh below MTL	-	more marsh below MTL represents more risk with SLR
	Shoreline	Hardened shoreline	-	more hardened shoreline represents more impediments to migration
Adaptive	Shoreline	Shoreline complexity	+	the more complex the shoreline the more protection it provides
Capacity	Migration Ability	Marsh migration space	igration space + more migration space represents a larger potential future area for mar	more migration space represents a larger potential future area for marsh to move into
	Tring action 7 tollicy	Wetland "connectedness"	+	marshes that are more connected in the future are more resilient





to sea level rise, and potential for adaptation.

Led by the National Estuarine Research Reserve System and the NOAA Office for Coastal Management, this study can help inform marsh restoration and conservation planning, identify marshes with particular characteristics across large areas, target fieldwork, and strengthen experimental design. It also can be used as a screening tool for national and regional work or be replicated with finer-scale data to create higher-resolution products.

Nationally Significant and Locally Relevant

The marsh resilience assessment was conducted with nationally standardized data sets for coastal areas of the lower 48 United States, not including the Great Lakes. The analysis systematically ranked resilience at multiple scales. It offers standardized comparisons of resilience across "marsh units" with broadly similar land use, land cover, and hydrology characteristics. Marsh resilience scores are summarized at the watershed scale (HUC-12 units).

Marsh Features

Marsh Data

Study Findings


Tidal marsh resilience to sea level rise was calculated using 13 GIS-based metrics related to current marsh conditions, vulnerability to sea level rise, and adaptive capacity. Values for each metric were scored based on a quantile distribution (10 categories), and then combined. Results were summarized to the watershed scale (HUC-12 units). The analysis includes all watersheds (from the National Watershed Boundary Dataset) that contained estuarine marsh (as derived from NOAA's Coastal Change Analysis Program 2016 land cover) in the contiguous United States (not including the Great Lakes).


One can view, analyze, and compare marsh resilience in three ways: overall resilience, resilience by category, or resilience by individual metric.

Overall Resilience Rankings

The map seen here highlights the overall score for each marsh unit.
Resilience scores range from 1 (red) for least resilient to 10 (green) for most resilient.


The study found that overall marsh resilience varies by region, with the Southeast having the most resilient marshes, and the Northeast having the least. Marshes were most resilient in Georgia and least resilient in New York. Different factors, explored in the study report, contributed to the resilience of each region's marshes.





Current Condition Factors

This resilience category includes metrics that represent characteristics of individual marsh features as well as land cover and land uses within the surrounding watershed. In this map you can use the buttons (below) to change the maps to see marsh unit rankings for each metric and more explanation of the metrics.






Vulnerability Factors

This resilience category includes metrics that represent possible current and future factors related to vulnerabilities of marsh features. In this map you can use the buttons (below) to change the maps to see marsh unit rankings for each metric and more explanation of the metrics.

Vulnerability Factor High/Low Q Columbus Category Rank This map highlights the overall vulnerability rating. This rating Low includes factors associated with the High erosion potential, tide range, and the current distribution of marshes within that tidal range. This information is related to these features as they exist today (potential future changes are not considered here). Chesapeak 6 Esri, GEBCO, DeLorme, NaturalVue | VGIN, Esri, HERE, Garmin, FAO, NOAA, USGS, EPA, NPS Powered by Esri Vulnerability Erodibility Tidal Range % Below MHHW % Below MTL




💠 webTA | Log In 🤌 E2 Travel 🔝 TOMIS | Task Order... 🔻 CCAPboard_JIRA 🔟 National Oceanic a... 🥞 OCM Intranet

Adaptive Capacity Factors

Adaptive capacity factors include metrics that represent characteristics of individual marshes that highlight the potential for that marsh to adapt to future sea level conditions. In this map you can use the buttons (below) to change the maps to see marsh unit rankings for each metric and more explanation of the metrics.

Adaptive Capacity Factor Category Rank High/Low Columbus This map highlights the overall High adaptive capacity rating. This rating includes factors associated with the Low shoreline configuration and hardened structures, as well as the space available for migration and the potential configuration of that future marsh area. Shoreline information is related to these features as they exist today (potential future changes are not considered here), while predictions of marsh migration are based on modeling done within NOAA Sea Level Rise Viewer. 6 Esri, GEBCO, DeLorme, NaturalVue I VGIN, Esri, HERE, Garmin, FAO, NOAA, USGS, EPA, NPS Powered by Esri Adaptive Capacity % Hardened Shoreline Shoreline Sinuosity Marsh Migration Space Future Marsh Connectedness





Applying the Data

This analysis and its findings can be used to advance understanding of the forces that drive tidal marsh resilience, and to support a range of efforts to manage and protect these precious resources for the future. Wetland managers can use data from this analysis to determine which marshes need help and what kind of help they may need.


💠 webTA | Log In 🤌 E2 Travel 🤡 TOMIS | Task Order... 🔻 CCAPboard_JIRA 🔟 National Oceanic a... 🤡 OCM Intranet


Data for Marsh Management	Marsh Resilience Category					
	Current condition	Vulnerability to sea level rise	Adaptive capacity	Interpretation		Management Options
Resilience rankings can inform where and how to invest in marsh management. This table	High	Low	High	A marsh in good condition that will migrate inland naturally over the long-term.	**	Land protection or policy: Fee or conservation easement purchase of marsh or migration space. Managed relocation of structures in marsh or migration space.
summarizes eight management options, each informed by how well the unit's marshes ranked in the	High	High	High	A marsh in good condition that will adapt if it can out-pace current vulnerability to sea level rise.	-	Adaptation (sediment): Increase sediment supply to current marsh footprint though strategies like thin layer sediment placement and removal of barriers to hydrologic flow within the marsh.
resilience categories (current conditions, vulnerability to relative sea level rise, and adaptive	High	Low	Low	A marsh in good condition that cannot maintain current footprint long-term without active management.		Adaptation (migration and land protection): Remove infrastructure or topographical barriers to inland migration. Protect marsh's current footprint.
capacity). Options are considered from an ecological, rather than	High	High	Low	A marsh in good condition but also vulnerable to sea level rise and cannot maintain current footprint without active management.		Phased adaptation: Implement thin layer sediment placement, but only if barriers to inland migration are removed.
socioeconomic, perspective. Note: Green text indicates a	Low	High	Low	A marsh in poor condition marsh that will not persist in the future.	-	Test innovative science or abandon marsh: Limit investment in land protection or restoration activities as effectiveness will be relatively short-term.
positive condition, red reflects a negative one. (Click the chart below to enlarge, or click here for a	Low	Low	Low	A marsh in poor condition not currently vulnerable to sea level rise, but cannot maintain current footprint without intervention.	-	Phased management (restore then adapt): Apply proven restoration techniques within current marsh footprint—e.g., invasive species management—but only if removal of barriers to migration are planned.
more detailed version of this chart.)	Low	High	High	A marsh in poor condition that has potential to migrate inland naturally, if it can out-pace sea level rise.		Phased management (restore then adapt): Apply proven restoration techniques within current marsh footprint that ideally also increase sediment supply to marsh, e.g. ditch remediation, tidal crossing restoration.
	Low	Low	High	A marsh in poor condition that is not currently vulnerable to sea level rise and has potential to migrate inland naturally.		Restoration: Decrease current stressors to marsh e.g., <i>Phragmittes</i> or invasive crab control. Mitigate water quality issues in surrounding watershed, e.g., reduce fertilizer application in residential and agricultural areas.






practitioners identify the appropriate techniques to use for restoration. For example, marshes with high vulnerability to sea level rise and strong potential for migration may be good candidates for experimental restoration techniques like thin layer placement. Esri, GEBCO, DeLorme, NaturalVue | VGIN, Esri, HERE, Garmin, FAO, NOAA, USGS, EPA







Coastal Change Analysis Program





Coastal Change Analysis Program (C-CAP)

- FGDC National Geospatial Data Asset
- Coastal expression of the NLCD
- Regional Products at 30-meter
 - Updated every 5 years (1996 2016)
 - Back as far as 1975 in some locations
 - 2021 update planned (2023 release)
- High Resolution Products at 1-meter
 - Historically expensive to produce
 - Faster, cheaper, better now possible
 - 2021 national buildout (by 2025)








Comparison of Resolutions

REGIONAL (30 METER)



LOCAL (1 METER)






High Resolution Land Cover

ORTHOIMAGE




LAND COVER





1-meter 2017 C-CAP


National Wetland Inventory





1-meter 2017 C-CAP

30-meter 2016 C-CAP





Upcoming High Resolution Mapping







Questions?

Nate Herold Physical Scientist nate.herold@noaa.gov

Visit us on the Digital Coast at:

coast.noaa.gov/digitalcoast/topics/coastal-land-cover.html

