



## Impact of Marine Heatwaves on Subsurface Hydrography and Dissolved Oxygen in the Chesapeake Bay

NATHAN SHUNK; PIERO L.F. MAZZINI; CASSIA PIANCA


## Marine Heat Waves (MHWs): Global Ocean


- Range in Intensity, Duration, and Spatial Scale
- Global Long-Term Trends
  - Increasing Frequency, Intensity, Duration, and Spatial Extent
- Drivers of SST Anomalies
  - Advection
  - Air-Sea Interactions







- MED 2003: Benthic Species die offs
- WA 2011: Sea Grass die off (Strydom et al., 2020)
- NEP 2015 & NWA 2012: Fishery Declines & Biogeography shifts (Cheung & Froelicher, 2020; Mills et al., 2013)
- Brazil 2020: Aggravate Estuarine Water Quality Issues (Brauko et al., 2020)







- **MED 2003:** Benthic Species die offs
- WA 2011: Sea Grass die off (Strydom et al., 2020)
- NEP 2015 & NWA 2012: Fishery Declines & Biogeography shifts (Cheung & Froelicher, 2020; Mills et al., 2013)
- Brazil 2020: Aggravate Estuarine Water Quality Issues (Brauko et al., 2020)







- MED 2003: Benthic Species die offs
- WA 2011: Sea Grass die off (Strydom et al., 2020)
- NEP 2015 & NWA 2012: Fishery Declines & Biogeography shifts (Cheung & Froelicher, 2020; Mills et al., 2013)
- Brazil 2020: Aggravate Estuarine Water Quality Issues (Brauko et al., 2020)






- MED 2003: Benthic Species die offs
- WA 2011: Sea Grass die off (Strydom et al., 2020)
- NEP 2015 & NWA 2012: Fishery Declines & Biogeography shifts (Cheung & Froelicher, 2020; Mills et al., 2013)
- Brazil 2020: Aggravate Estuarine Water
   Quality Issues (Brauko et al., 2020)






- MED 2003: Benthic Species die offs
- WA 2011: Sea Grass die off (Strydom et al., 2020)
- NEP 2015 & NWA 2012: Fishery Declines &
   Biogeography shifts (Cheung & Froelicher, 2020;
   Mills et al., 2013)
- Brazil 2020: Aggravates Estuarine Water Quality Issues (Brauko et al., 2020)




- Anomalously warm: 90<sup>th</sup> percentile above climatology
- Prolonged:Period of at least 5 days
- **Discrete**:

  2-day gap between two 5-day intervals




- Anomalously warm:
  90<sup>th</sup> percentile above climatology
- Prolonged:Period of at least 5 days
- **Discrete**:

  2-day gap between two 5-day intervals



- Anomalously warm:
   90<sup>th</sup> percentile above climatology
- Prolonged:
   Period of at least 5 days
- **Discrete**:


  2-day gap between two 5-day intervals



- Anomalously warm:

   90<sup>th</sup> percentile above
   climatology
- Prolonged:
   Period of at least 5 days
- **Discrete**:


  2-day gap between two 5-day intervals



- Anomalously warm:


   90<sup>th</sup> percentile above
   climatology
- Prolonged:Period of at least 5 days
- Discrete:

  2-day gap between two 5-day intervals



# Marine Heatwaves in the Bay

- Increasing Frequency of Events
- Greater Percentage in Summer
- Increasing Cumulative Intensity
- Co-occurrence Across the Bay



Mazzini & Pianca, 2022

### Research Objectives

Characterize the **spatial distribution** of MHWs along the Chesapeake Bay estuarine salinity gradient, and their **seasonal variability**.

Investigate **anomalies** in hydrographic and water quality parameters associated with MHWs, specifically **salinity** and **dissolved oxygen** content.

Analyze how mixed layer depth and vertical stratification may impact distribution of temperature anomalies in the water column during MHW events.


- **Data:** 21 Stations; 1986-2021
  - Continuous Monitoring Stations (NOAA, NERR, VIMS)

(Mazzini & Pianca, 2022)

- Chesapeake Bay Program Cruise Profiles
  - Temperature, Salinity, Dissolved Oxygen
  - Density Calculated Using Gibbs-SeaWater Toolbox TEOS-10

(McDougall & Baker, 2011)

• Profiles Linearly Interpolated




- Data: 21 Stations; 1986-2021
  - Continuous Monitoring Stations (NOAA, NERR, VIMS)


(Mazzini & Pianca, 2022)


- Chesapeake Bay Program Cruise Profiles
  - Temperature, Salinity, Dissolved Oxygen
  - Density Calculated Using Gibbs-SeaWater Toolbox TEOS-10

(McDougall & Baker, 2011)


- Profiles Linearly Interpolated
- MHW Identification
  - Spatially Averaged Time Series
  - m\_mhw (Zhao & Marin, 2019)
  - Climatology Not Smoothed
  - 5 Day Duration









#### MHW Profiles per station per month






- Daily Climatology at each grid point
- Produce Daily Anomaly profiles




$$y(t) = \mu + \sum \beta_m \sin(m\pi f_0 t) + \beta_m \cos(m\pi f_0 t)$$
$$f_0 = 1/365.2422; \ m = 2,4,6$$

- Daily Climatology at each grid point
- Produce Daily Anomaly profiles
- LSM Harmonic Analysis: 3 Harmonics




$$y(t) = \mu + \sum \beta_m \sin(m\pi f_0 t) + \beta_m \cos(m\pi f_0 t)$$
$$f_0 = 1/365.2422; \ m = 2,4,6$$

- Daily Climatology at each grid point
- Produce Daily Anomaly profiles
- LSM Harmonic Analysis: 3 Harmonics




$$y(t) = \mu + \sum \beta_m \sin(m\pi f_0 t) + \beta_m \cos(m\pi f_0 t)$$
$$f_0 = 1/365.2422; \ m = 2,4,6$$

- Daily Climatology at each grid point
- Produce Daily Anomaly profiles
- LSM Harmonic Analysis: 3 Harmonics
- Validated against Monthly Climatology



$$y(t) = \mu + \sum \beta_m \sin(m\pi f_0 t) + \beta_m \cos(m\pi f_0 t)$$
$$f_0 = 1/365.2422; \ m = 2,4,6$$

- Daily Climatology at each grid point
- Produce Daily Anomaly profiles
- LSM Harmonic Analysis: 3 Harmonics
- Validated against Monthly Climatology




$$y(t) = \mu + \sum \beta_m \sin(m\pi f_0 t) + \beta_m \cos(m\pi f_0 t)$$
$$f_0 = 1/365.2422; \ m = 2,4,6$$

#### For Temperature & Salinity

- Daily Climatology at each grid point
- Produce Daily Anomaly profiles
- LSM Harmonic Analysis: 3 Harmonics
- Validated against Monthly Climatology


#### For Dissolved Oxygen

Monthly Average Climatology





### Results: Monthly Mean MHW Temperature Anomalies


Mixed Layer Depth
Used Irby et al. (2016)
Definition





#### Results: Monthly Mean MHW Temperature Anomalies












### Summary: Temperature and DO Anomalies



Contact:

Nathan Shunk- <a href="mailto:npshunk@vims.edu">npshunk@vims.edu</a>
Piero Mazzini- <a href="mailto:pmazzini@vims.edu">pmazzini@vims.edu</a>

#### Conclusions

- No Discernible Patterns in Salinity Anomalies
- Winter Temperature Anomalies Larger and Penetrate at Depth
- <u>Summer</u> Temperature Anomalies **Smaller** and <u>do not</u> **Penetrate at Depth**

Contact:

Nathan Shunk- <a href="mailto:npshunk@vims.edu">npshunk@vims.edu</a>
Piero Mazzini- <a href="mailto:pmazzini@vims.edu">pmazzini@vims.edu</a>

#### Conclusions

#### For Temperature & Salinity

- No Discernible Patterns in Salinity Anomalies
- Winter Temperature Anomalies Larger and Penetrate at Depth
- <u>Summer</u> Temperature Anomalies Smaller and do not Penetrate at Depth

#### For Dissolved Oxygen

- Winter DO anomalies are negative throughout water column
- Summer DO anomalies are negative in the middle of the water column Capped by Surface Mixed Layer

Contact:

Nathan Shunk- <a href="mailto:npshunk@vims.edu">npshunk@vims.edu</a>
Piero Mazzini- <a href="mailto:pmazzini@vims.edu">pmazzini@vims.edu</a>

#### Conclusions

#### For Temperature & Salinity

- No Discernible Patterns in Salinity Anomalies
- Winter Temperature Anomalies Larger and Penetrate at Depth
- <u>Summer Temperature Anomalies Smaller and do not Penetrate at Depth</u>


#### For Dissolved Oxygen

- Winter DO anomalies are negative throughout water column
- Summer DO anomalies are negative in the middle of the water column Capped by Surface Mixed Layer

#### **Future Work**

- Examine Chlorophyll Anomalies
- Investigate MLD Definition and Temperature Anomaly Discrepancies

### Appendix: Salinity



### Appendix: MHW Events

