

# High-resolution Land Use and Hydrography Data to Inform the Phase 7 Watershed Model

Peter R. Claggett<sup>1</sup>, Labeeb Ahmed<sup>1</sup>, Matthew Baker<sup>4</sup>, Jacob Czawlytco<sup>2</sup>, Sean MacFaden<sup>3</sup>, Sarah McDonald<sup>1</sup>, Jarlath O'Neil-Dunne<sup>3</sup>, David Saavedra<sup>2</sup>, and Rachel Soobitsky<sup>2</sup>

- 1 Presenting Author, Lower Mississippi-Gulf Water Science Center, U.S. Geological Survey, Annapolis, MD 21403
- 2 Chesapeake Conservancy, Annapolis, MD 21403
- 3 University of Vermont Spatial Analysis Laboratory, Burlington, VT 05405
- 4 University of Maryland, Baltimore County, Baltimore, MD 21250

## Modeling Workgroup Quarterly Meeting April 6, 2021

### **Disclaimer**

"This information is preliminary or provisional and is subject to revision. It is being provided to meet the need for timely best science. The information has not received final approval by the U.S. Geological Survey (USGS) and is provided on the condition that neither the USGS nor the U.S. Government shall be held liable for any damages resulting from the authorized or unauthorized use of the information."



### **One-meter Resolution Products Relevant to Phase 7 Model**

Land Use: 2013, 2017, and 2021; 60+ classes; accurate change over time.

2017-18 data highlights: addition of lakes/ponds, solar fields, and orchards/vineyards; improved mapping of tidal and non-tidal wetlands

Planned public release date: December 2021

2021-22 data highlights: addition of open channel, tree canopy over channel, and animal operations.

Planned public release date: June 2023

### Hydrography: 2-D raster (discontinuous), 1-D polyline (continuous)

Features: streams, rills/gullies, agricultural ditches, roadside ditches, floodplains (from FACET), floodplain depressions, detention features, headwater wetlands, other.

Attributes: width, bank height, flow permanence, stream order, drainage area

Planned public release date: June 2024 (draft products will be available in 2022-23)

#### **CBP 2017 Land Use Classification (27 macro-classes, version 1)**

#### 1. Water (3)

1.1 Lentic

1.1.1 Estuary (tidal)

1.1.2 Lakes & Ponds

1.2 Lotic

1.2.1 Streams

#### 2. Developed (8)

2.1 Impervious

2.1.1 Roads

2.1.2 Structures

2.1.3 Other Impervious (Parking lots, driveways)

2.1.4 Tree Canopy (TC) over Impervious

#### 2.2 Pervious

2.2.1 Turf Grass

2.2.2 Bare Developed

2.2.3 Suspended Succession (rights-of-way)

2.2.4 Tree Canopy over Turf Grass

#### 3. Forest (4)

3.1 Forest (>= 1 acre, 240-ft width)

3.2 Tree Canopy in Agriculture

3.3 Harvested Forest (<= 3 years)

3.4 Natural Succession (> 3 years)

#### 4. Production (7)

4.1 Agriculture

4.1.1 Cropland

4.1.2 Pasture

4.1.3 Orchard/vineyard

4.2 Solar fields

4.2.1 Impervious

4.2.2 Pervious

4.3 Extractive (active mines)

#### 5. Wetlands and Water Margins (5)

5.1 Tidal

5.2 Riverine (Non-tidal)

5.2.1 Headwater

5.2.2 Floodplain

5.3 Terrene/Isolated (Non-tidal)

5.4 Bare shore


### **CBP Final Land Use Classification (64 classes)**


1 Water (9)	. Forest (7)	
1. Water (0)	3.1 Forest (>= 1 acre, 240-ft width)	4.3 Extractive (active mines)
1.1 Lentic	3.2 Tree Canopy in Agriculture	4.3.1 Barren
1.1.1 Estuary (tidal)	3.3 Harvested Forest (<= 3 years)	4.3.2 Impervious
1.1.2 Lakes & Ponds	3.3.1 Barren	
1.2 Lotic	3.3.2 Herbaceous	5. Wetlands and Water Margins (21
1.2.1 Streams	3.4 Natural Succession (> 3 years)	5.1 Tidal
1.2.1.1 Open Channel	3.4.1 Barren	5.1.1 Barren
1.2.1.2 Tree Canopy over Channel	3.4.2 Herbaceous	5.1.2 Herbaceous
1.2.1.3 Culverted/ Buried Channel	3.4.3 Scrub-shrub	5.1.3 Scrub-shrub
1.2.2.Ditches	0.4.0 Colds 3111ds	5.1.4 Tree Canopy
1.2.2.1 Open Ditch	. Production (16)	5.1.5 Forest
1.2.2.2 Tree Canopy over Ditch 1.2.2.3 Culverted/ Buried Ditch		5.2 Riverine (Non-tidal)
	4.1 Agriculture	5.2.1 Headwater
	4.1.1 Cropland	5.2.1.1 Barren
2. Developed (12)	4.1.1.1 Barren	5.2.1.2 Herbaceous
2.1 Impervious	4.1.1.2 Herbaceous	5.2.1.3 Scrub-shrub
2.1.1 Roads	4.1.2 Pasture	5.2.1.4 Tree Canopy
2.1.2 Structures	4.1.2.1 Barren	5.2.1.4 Tree Carlopy 5.2.1.5 Forest
2.1.3 Other Impervious (Parking lots, driveways)	4.1.2.2 Herbaceous	
2.1.4 Tree Canopy (TC) over Impervious	4.1.3 Orchard/vineyard	5.2.2 Floodplain
2.1.4.1 TC over Roads	4.1.3.1 Barren	5.2.2.1 Barren
2.1.4.2 TC over Structures	4.1.3.2 Herbaceous	5.2.2.2 Herbaceous
2.1.4.3 TC over Other Impervious	4.1.3.3 Scrub-shrub	5.2.2.3 Scrub-shrub
2.2 Pervious	4.1.4 Animal Operations (TBD)	5.2.2.4 Tree Canopy
2.2.1 Turf Grass	4.1.4.1 Impervious	5.2.2.5 Forest
2.2.2 Bare Developed	4.1.4.2 Barren	5.3 Terrene/Isolated (Non-tidal)
2.2.3 Suspended Succession (rights-of-way)	4.1.4.3 Herbaceous	5.3.1 Barren
2.2.3.1 Barren		5.3.2 Herbaceous
2.2.3.2 Herbaceous	4.2 Solar fields	5.3.3 Scrub-shrub
2.2.3.3 Scrub-shrub	4.2.1 Impervious	5.3.4 Tree Canopy
2.2.4 Tree Canopy over Turf Grass	4.2.2 Pervious	5.3.5 Forest
2.2.4 free Sanopy over full Stass	4.2.2.1 Barren	5.4 Bare shore
	4.0.0.0.1.1	


4.2.2.3 Scrub-shrub

### Parcel Segments: the key to classifying high-resolution land use











Finer-scale representation of sources and sinks of TN, TP, and TSS.




### **Comparing the Census of Agriculture and High-res Land Use**


		Census	Мар
FIPS	CNTY_NAME	AG_17	AG_17
10005	SUSSEX	238,450	234,831
24005	BALTIMORE	56,655	60,696
24035	QUEEN ANNES	131,656	118,952
24045	WICOMICO	61,747	59,732
36017	CHENANGO	82,944	109,915
42015	BRADFORD	167,068	231,889
42033	CLEARFIELD	28,875	104,169
42041	CUMBERLAND	141,157	116,748
42071	LANCASTER	325,806	304,277
51015	AUGUSTA	213,135	145,181
51073	GLOUCESTER	14,370	13,027
51107	LOUDOUN	83,858	55,791
54003	BERKELEY	44,859	32,009
54031	HARDY	76,819	37,251









### **Tidal Wetlands: Maryland and Delaware Wetland Datasets**




So what?

Increased area of tidal wetlands = decreased land area in Watershed Model























Accurate characterization of the nutrient loading implications of land use change requires understanding land use history: 1980's to present.





# Parcel-Level Assessment of Historical Land Use: 1985 - 2017

Year-Built Attributes from Tax Records



yearblt

1985-1989

1990-1994

1995-1999

2000-2004

2005-2009

2010-2014


2015-2017

Year-Built Attributes from USGS' LCMAP



### **Hyper-Resolution Hydrography**


- 1. Lidar elevation
- 2. Valley-scale geomorphons
- 3. Channel-scale geomorphons
- 4. Extract valley network
- 5. Extract channels using valley network
- 6. QAQC channel skeleton
- 7. Connect stream network




### **Alternative Stream Resolutions**

Lower Susquehanna Example


National Hydrography Dataset, 1:24,000 6,923.6 km



CBP Hyper-Resolution Streams, 1:2000 16,784.6 km




### Why the 2x difference in stream length?



So what?

- Increased travel time within concentrated flow paths
- Decreased travel time over/through land

What are the nutrient/sediment transport and transformation implications of these data? Will Phase 7 be sensitive to these changes?



### Floodplain Mapping (FACET)

Patuxent River @ HW 50

**Step 1:** Identify active floodplain extent in field based on topography, vegetation, evidence of recent flooding (e.g. fine sediment deposits, debris deposits aligned perpendicular to the channel), and flood stage magnitudes at nearby stream gages

**Step 2:** Identify HAND threshold aligning with field-measured floodplain extent at each field site

**Step 3:** Predictive linear model related HAND height thresholds to drainage area and physiographic province

