

Old "News":

Forested vs. Grass Buffers

Proc. of Nat'l Acad of Science, Sept 2004

- 16 stream segments in SE PA
- Flowing in and out of woodlots, pastures
- Findings for forested vs. grass buffers:
 - 2 to 9x nitrogen uptake in forested reaches
 - 3-5x more total biological activitiy
 - 2-3x wetted channel width

Literature review on buffer width:

- Reviewed all literature on buffer width
- Conclusion: science supports a 100' width as a defensible goal
- Width's impact varies with parameter
 - Temp. effects gotten with narrower buffers
 - -Sediment:
 - first 10m stops majority by weight (sand)
 - next 20m stops smaller portion (silt/clay)

Hurricane Sandy Project:

- Headwaters catchment ~4 sq mi
- Majority of landowners doing:
 - –level lip spreaders
 - -forested buffers 100' per side
- A few doing:
 - -woody debris, created wetlands
- All overlaid with science









Pre-emergence herbicide:

- Bird's deposit seeds of invasive spp.
- Snapshot TM being tested INSIDE tubes

Battling reed canary grass:

- Prep mowing before planting
- Routine herbicide spots (2x/yr)
- Routine follow-up mowing
- So far, success

Battling multiflora rose + mile-a-minute:

- Prep mowing before planting (\$2k/ac)
- Planted as usual with shelters
- Broadcast herbicide for MFR, MAM
- Reseeded with pasture mix
- Weeded INSIDE tubes
- Now into routine maintenance (mowing + herbicide spots)





Stay tuned for findings on:

- herbicide spot size
- initial plant size: 3x3x9" vs. 2 gallon etc.
- tube tests: Plantra TM, Shelterguard TM
- fiberboard mat w/cherry extract
- 4' welded wire fence to protect shrub clusters (biodegrad. version: wooden stakes + twine)







Contact Info:

David Wise
Stroud Water
Research Center
717-979-9997
dwise@stroudcenter.org

