

# Why Tree Canopy Land Uses in Phase 6?

- Since 2003, it has been the policy of the Chesapeake Bay Program partners to increase urban tree canopy cover for water quality and other benefits
  - Reaffirmed and strengthened in the 2014 Chesapeake Bay Agreement Tree Canopy Outcome
- Urban tree canopy benefits are not directly accounted for in the CB Model land uses
  - Implication: Retaining tree canopy has no “value” in the TMDL framework, whereas Forest, Open Space, and other preferred land uses do



# Tree Canopy LU/Loading Rate Review Timeline

- **February 11** – Webinar, comments due 2/22
- Brief workgroups and seek their approval of Tree Canopy Land Uses/Loading rates where needed
  - **March 2:** Forestry Workgroup
  - **March 2:** Land Use Workgroup
  - **March 3:** Watershed Technical Workgroup
  - **March 8:** Urban Stormwater Workgroup
  - **March 10:** Modeling Workgroup
  - **March 14:** Water Quality Goal Implementation Team
- **Goal to include Tree Canopy land uses/loading rates in April 1 beta calibration**
- Expert Panel report on Tree Canopy BMP for new plantings – partnership review starting in April

# Relative Non-Point Source Pollution Loading Rates of Tree Canopy Land Uses



**Justin Hynicka<sup>1</sup> and Marion Divers<sup>2</sup>**

**1 – Maryland Department of Natural Resources – Forest Service**

**2 – University of Pittsburgh – Department of Geology and Environmental Science**





## This analysis builds on work by the Tree Canopy EP

---

- Expands the scope of the previous Tree Canopy Expert Panel Technical Memo beyond canopy interception to include ecosystem level processes
- This analysis draws from an expanded literature review on urban tree planting and canopy (Karen Cappiella, Center for Watershed Protection)
- Describe how changes in water yield due to tree canopy interact with nutrient cycling processes to reduce non-point source pollution
- **Incorporates feedback from webinar on 02/11/16**

# Why water balance, and what does it look like?


- Regardless of the source, nutrients and sediment are transported by water to streams, rivers, estuaries, and beyond.
- Water balance is a mathematical framework that accounts for all additions and losses of water from a drainage area



# Runoff calculated using the SCS Curve Number Method

- Developed by the USDA Soil Conservation Service (TR-55, 1986)
- Added a term to account for tree canopy interception ( $C_i$ ), which isolates the effects of tree canopy from the water retaining properties of the underlying land use.
- $C_i$  ranges from 0.02 to 0.11 inches of precipitation per storm for deciduous tree species, and 0.02 to 0.18 in. per storm for coniferous trees (Breuer et al. 2003).
- We used a fixed  $C_i$  value of 0.05 in. in our calculations during the growing season (April through October).
- $C_i$  set to zero in the winter.**


$$R = \frac{(P - C_i - I_a)^2}{(P - C_i - I_a) + S}$$



# Modified CN assumptions to better reflect runoff at small scales


- Observations at that the watershed scale during CN method development revealed that  $I_a/S \sim 0.2$  (Garen and Moore 2005)
- More recent work has revealed that this simplification underestimates runoff of small storm events especially at smaller scales (Woodward et al. 2003)
- $I_a/S \sim 0.05$  is more appropriate for evaluating the role of tree canopy in runoff calculations (Woodward et al. 2003)
- In addition, soils from hydrologic soil group C (rather than D) are likely more representative conditions in the urban environment.

$$R = \frac{(P - C_i - I_a)^2}{(P - C_i - I_a) + S}$$







# We used daily weather data (2005 to 2015) from eight locations



— Chesapeake Bay Watershed Boundary




(NCDC Quality Controlled Local Climatological Data)





# Leaching calculated by tracking changes in soil water volume


- Leaching is water that infiltrates in excess of the soil water holding capacity, which varies by soil type and over time due to plant evapotranspiration (ET).
- Calculations based on silt clay loam soils with a maximum water holding capacity of 2.0 inches per foot of soil (Brady and Weil 1996)



(<http://soilquality.org.au/factsheets/water-availability>)


# Leaching calculated by tracking changes in soil water volume

- Leaching is water that infiltrates in excess of the soil water holding capacity, which varies by soil type and over time due to plant evapotranspiration (ET).
- Average annual ET is similar between grasses, natural forests, and urban trees ranging from  $\sim 15$  to  $24 \text{ in yr}^{-1}$  (Ford 2011, Penmen 1948, Wullschleger 2001, Wullschleger 2000, Wilson 2001, and Peters 2010)
- ET during growing was set at 0.05 and 0.08 inches per day for turfgrass and canopy over turfgrass, respectively
- **During the dormant season these land uses are equivalent**



# Throughflow provides a source of water and nutrients to trees

- Average daily throughflow was estimated using the volume of water leached annually from a square meter of turfgrass and redistributing it evenly over the course of a year.
- ET of trees during growing season set to 0.05 inches per day (Ford 2011, Wullschleger 2001, Wullschleger 2000, Wilson 2001, and Peters 2010)



# Updated water yield results

---

Land Use	Precip. (in)	Runoff Red. (%)	Leaching Red. (%)	Throughflow Red. (%)	Total (%)
Canopy over Turfgrass	39.9	29.0	22.5	NA	23.8
Canopy over Impervious	39.9	7.0	NA	22.3	14.9

# Updated water yield results

---

Land Use	Precip. (in)	Runoff Red. (%)	Leaching Red. (%)	Throughflow Red. (%)	Total (%)
Canopy over Turfgrass	39.9	29.0	22.5	NA	23.8
Canopy over Impervious	39.9	7.0	NA	22.3	14.9

# Hydrologic processes govern the fate and transport of pollution

---

- Absolute loading rates for TC land uses are limited by the low availability of concentration data

$$J_{gc} = \overline{X}_1 \cdot \sum R_{gc} + \overline{X}_2 \cdot \sum L_{gc}$$



# Hydrologic processes govern the fate and transport of pollution

- Absolute loading rates for TC land uses are limited by the low availability of concentration data



# Hydrologic processes govern the fate and transport of pollution

---

- Absolute loading rates for TC land uses are limited by the low availability of concentration data

$$\frac{J_{gc}}{J_g} = \frac{\overline{X}_1 \cdot \sum R_{gc} + \overline{X}_2 \cdot \sum L_{gc}}{\overline{X}_3 \cdot \sum R_g + \overline{X}_4 \cdot \sum L_g}$$

- For a long-term practice in complex watersheds modeling is the best approach to estimate relative loading rates among land classes
- If trees reduce edge of field pollution fluxes then where does the mitigated pollution go?

# Trees promote pollution storage and removal in pervious areas

- N and P are essential nutrients that are taken up through roots and stored in plant tissues
- Trees increase infiltration rates that leads to greater filtration/capture of nutrients and sediments (Bartens 2008, Busman 2002, Day 2010, Leguedois 2008)
- Increased soil moisture and soil organic matter from trees enhances the conditions required for denitrification (Day 2010, Gift 2010, Huyler 2014, Lovett 2002, Takahashi 2008, Zhu 2004)



# Impervious surfaces limit EOF water quality benefits of trees

- New N and P inputs of have little chance to enter the nutrient cycle
- Our estimate of throughflow is poorly constrained, and a large portion of pollution taken up by trees with canopy over impervious is later returned to that surface.
- Recommended relative loading rate of 7% for N, P, and sediment was based solely on downstream benefits of reduced runoff (Asadian 2009, Nowak 2007, Wang 2008)



# Impervious surfaces limit EOF water quality benefits of trees

---

- New N and P inputs of have little chance to enter the nutrient cycle
- Our estimate of throughflow is poorly constrained, and a large portion of pollution taken up by trees with canopy over impervious is later returned to that surface.
- Recommended relative loading rate of 7% for N, P, and sediment was based solely on downstream benefits of reduced runoff (Asadian 2009, Nowak 2007, Wang 2008)
- **However, N and P in wood is a long term store of water quality pollution**



# Estimating N and P stored in woody biomass

---

- We used estimates of annual wood production (Nowak and Crane 2002), annual leaf litter production (Abelho 2001, Chapin 2011, Martin 1998, and Olsen 1963), and annual fluxes of N and P deposition (NADP, and Smullen 1982)
- Combined fluxes with nutrient concentration data (Martin 1998, McGroddy 2004, Petterson 1984, Rastetter 1991)
- **What proportion of N and P stored in wood relative to total nutrient uptake (wood + leaf production) and new inputs of atmospheric deposition?**



# Estimating N and P stored in woody biomass



- Proportion of annual N and P stored in wood ~ 5 and 14 %, respectively
- We still need to make an assumption about uptake efficiency to make results comparable to water yield results.
- Assumed uptake efficiency based on the proportion of time that deciduous trees transpire water  $7/12 \text{ months} \times 1/2 \text{ hours/day} = 0.29$

## Final estimated relative loading rates for TC land uses

---

Land Use	Total N Reduction (%)	Total P Reduction (%)	Sediment Reduction (%)
Canopy over Turfgrass	23.8	23.8	29.0 x 0.2
Canopy over Impervious	7.0 + 1.5	7.0 + 4.0	7.0

# Final estimated relative loading rates for TC land uses

Land Use	Total N Reduction (%)	Total P Reduction (%)	Sediment Reduction (%)
Canopy over Turfgrass	23.8	23.8	5.8
Canopy over Impervious	8.5	11.0	7.0

Relative loading rates from 2/11/16 webinar

Land Use	Pollution Reduction (%)
Canopy over Turfgrass	26.0
Canopy over Impervious	7.1

# Final estimated relative loading rates for TC land uses

Land Use	Total N Reduction (%)	Total P Reduction (%)	Sediment Reduction (%)
Canopy over Turfgrass	23.8	23.8	5.8
Canopy over Impervious	8.5	11.0	7.0

